

EV

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number  
WO 02/32893 A2

(51) International Patent Classification: C07D 401/00

(21) International Application Number: PCT/US01/32151

(22) International Filing Date: 15 October 2001 (15.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/240,901 17 October 2000 (17.10.2000) US

(71) Applicant: SCHERING CORPORATION [US/US];  
Patent Department - K-6-1 1990, 2000 Galloping Hill  
Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors: ASLANIAN, Robert, G.; 144 Philip Drive,  
Rockaway, NJ 07866 (US). SHIH, Neng-Yang; 1 Maple  
Drive, North Caldwell, NJ 07006 (US). TING, Pauline,  
C.; 74 Delwick Lane, New Providence, NJ 07974 (US).  
BERLIN, Michael, Y.; 14 Hendrick Road, Flemington,  
NJ 08822 (US). ROSENBLUM, Stuart, B.; 16 Steven  
Terrace, West Orange, NJ 07052 (US). MC CORMICK,  
Kevin, D.; 5 Pace Drive, Edison, NJ 08820 (US). TOM,  
Wing, C.; 133 Cedar Grove Parkway, Cedar Grove, NJ  
07009 (US). BOYCE, Christopher, W.; 71 Bonnell Street,  
Flemington, NJ 08822 (US). MANGIARACINA, Pietro;  
4 Montclair Avenue, Monsey, NY 10952 (US). MUTAHL,

Mwangi, Wa; 45 Snyder Road, Fords, NJ 08863 (US). PI-  
WINSKI, John, J.; 6 Saddle Ridge Drive, Clinton Town-  
ship, NJ 08833 (US).

(74) Agent: JEANETTE, Henry, C.; Schering-Plough Corpo-  
ration, Patent Department - K-6-1 1990, 2000 Galloping  
Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,  
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,  
ID, IL, IN, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LU,  
LV, MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL,  
PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,  
UZ, VN, YU, ZA.


(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL NON-IMIDAZOLE COMPOUNDS



(57) Abstract: Disclosed are novel compounds of  
the formula: (I) Also disclosed are pharmaceutical  
compositions comprising the compounds of For-  
mula (I). Also disclosed are methods of treating  
various diseases or conditions, such as, for exam-  
ple, allergy, allergy-induced airway responses, and  
congestion (e.g., nasal congestion) using the com-  
pounds of Formula (I). Also disclosed are methods  
of treating various diseases or conditions, such as,

for example, allergy, allergy-induced airway responses, and congestion (e.g., nasal congestion) using the compounds of Formula (I)  
in combination with a H1 receptor antagonist.

WO 02/32893 A2

5

## NOVEL NON-IMIDAZOLE COMPOUNDS

### 10 REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application Serial No. 60/240901 filed October 17, 2000.

### BACKGROUND OF THE INVENTION

15 WO 95/14007 published May 26, 1995 discloses H<sub>3</sub> receptor antagonists of the imidazole type.

WO99/24405 published May 20, 1999 discloses H<sub>3</sub> receptor ligands of the imidazole type.


20 US 5,869,479 issued February 9, 1999 discloses compositions for the treatment of the symptoms of allergic rhinitis using a combination of at least one histamine H<sub>1</sub> receptor antagonist and at least one histamine H<sub>3</sub> receptor antagonist.

In view of the art's interest in compounds which affect H<sub>3</sub> receptors, novel compounds that are antagonists of H<sub>3</sub> receptors would be a welcome contribution to the art. This invention provides just such a contribution.

25

### Summary of the Invention

The present invention provides novel compounds of structure I.



30 or a pharmaceutically acceptable salt or solvate thereof, wherein:

(1) R<sup>1</sup> is selected from:


- 2 -

- (a) aryl;  
 (b) heteroaryl;  
 (c) heterocycloalkyl  
 (d) alkyl;  
 (e) cycloalkyl; or  
 (f) alkylaryl;

wherein said  $R^1$  groups are optionally substituted with 1 to 4 substituents independently selected from:

- (1) halogen (e.g., Br, F, or Cl, preferably F or Cl);  
 (2) hydroxyl (i.e., -OH);  
 (3) lower alkoxy (e.g.,  $C_1$  to  $C_6$  alkoxy, preferably  $C_1$  to  $C_4$  alkoxy, most preferably  $C_1$  to  $C_2$  alkoxy, more preferably methoxy);  
 (4)  $-CF_3$ ;  
 (5)  $CF_3O-$ ;  
 (6)  $-NR^4R^5$ ;  
 (7) phenyl;  
 (8)  $-NO_2$ ;  
 (9)  $-CO_2R^4$ ;  
 (10)  $-CON(R^4)_2$  wherein each  $R^4$  is the same or different;  
 (11)  $-S(O)_mN(R^{20})_2$  wherein each  $R^{20}$  is the same or different H or alkyl group, preferably  $C_1$  to  $C_4$  alkyl, most preferably  $C_1$ - $C_2$  alkyl, and more preferably methyl;  
 (12)  $-CN$ ; or  
 (13) alkyl; or

- (2)  $R^1$  and X taken together form a group selected from:



- (3) X is selected from:  $=C(O)$ ,  $=C(NOR^3)$ ,  $=C(NNR^4R^5)$ ,



- 3 -

- (4)  $M^1$  is carbon;  
 (5)  $M^2$  is selected from C or N;  
 (6)  $M^3$  and  $M^4$  are independently selected from C or N;  
 (7) Y is selected from: is  $-CH_2-$ ,  $=C(O)$ ,  $=C(NOR^{20})$  (wherein  $R^{20}$  is as

5 defined above), or  $=C(S)$ ;

- (8) Z is a  $C_1 - C_6$  alkyl group;

(9)  $R^2$  is a five or six-membered heteroaryl ring, said six-membered heteroaryl ring comprising 1 or 2 nitrogen atoms with the remaining ring atoms being carbon, and said five-membered heteroaryl ring containing 1 or 2 heteroatoms  
 10 selected from: nitrogen, oxygen, or sulfur with the remaining ring atoms being carbon; said five or six membered heteroaryl rings being optionally substituted with 1 to 3 substituents independently selected from: halogen, hydroxyl, lower alkyl, lower alkoxy,  $-CF_3$ ,  $CF_3O-$ ,  $-NR^4R^5$ , phenyl,  $-NO_2$ ,  $-CO_2R^4$ ,  $-CON(R^4)_2$  wherein each  $R^4$  is the same or different,  $-CH_2NR^4R^5$ ,  $-(N)C(NR^4R^5)_2$ , or  $-CN$ ;


- 15 (10)  $R^3$  is selected from:

- (a) hydrogen;  
 (b)  $C_1 - C_6$  alkyl;  
 (c) aryl;  
 (d) heteroaryl;  
 20 (e) heterocycloalkyl;  
 (f) arylalkyl (e.g., aryl( $C_1$  to  $C_4$ )alkyl, e.g.,  $-(CH_2)_w$ aryl wherein w is 1 to 4, preferably 1 or 2, and most preferably 1, such as, for example  $-CH_2$ phenyl or  $-CH_2$ substituted phenyl);

- (g)  $-(CH_2)_e-C(O)N(R^4)_2$  wherein each  $R^4$  is the same or different,

- 25 (h)  $-(CH_2)_e-C(O)OR^4$ ;

- (i)  $-(CH_2)_e-C(O)R^{30}$  wherein  $R^{30}$  is a heterocycloalkyl group, such as, for example, morpholinyl, piperidinyl, piperazinyl or pyrrolidinyl, including



- (j)  $-CF_3$ ; or

- 30 (k)  $-CH_2CF_3$ ;

- 4 -

wherein said aryl, heteroaryl, heterocycloalkyl, and the aryl portion of said arylalkyl are optionally substituted with 1 to 3 (preferably 1) substituents selected from:

halogen (e.g., F or Cl), -OH, -OCF<sub>3</sub>, -CF<sub>3</sub>, -CN, -N(R<sup>45</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>45</sup>, or -C(O)N(R<sup>45</sup>)<sub>2</sub>, wherein each R<sup>45</sup> is independently selected from: H, alkyl, alkylaryl, or alkylaryl

5 wherein said aryl moiety is substituted with 1 to 3 substituents independently selected from -CF<sub>3</sub>, -OH, halogen, alkyl, -NO<sub>2</sub>, or -CN;

(11) R<sup>4</sup> is selected from: hydrogen, C<sub>1</sub> - C<sub>6</sub> alkyl, aryl, alkylaryl, said aryl and alkylaryl groups being optionally substituted with 1 to 3 substituents selected from: halogen, -CF<sub>3</sub>, -OCF<sub>3</sub>, -OH, -N(R<sup>45</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>45</sup>, -C(O)N(R<sup>45</sup>)<sub>2</sub>, or -CN; wherein R<sup>45</sup> is  
10 as defined above;

(12) R<sup>5</sup> is selected from: hydrogen, C<sub>1</sub> - C<sub>6</sub> alkyl, -C(O)R<sup>4</sup>, -C(O)<sub>2</sub>R<sup>4</sup>, or -C(O)N(R<sup>4</sup>)<sub>2</sub> wherein each R<sup>4</sup> is independently selected, and R<sup>4</sup> is as defined above;

(13) or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are bound forms a five or six membered heterocycloalkyl ring (e.g., morpholine);

15 (14) R<sup>6</sup> is selected from: alkyl, aryl, alkylaryl, halogen, hydroxyl, lower alkoxy, -CF<sub>3</sub>, CF<sub>3</sub>O-, -NR<sup>4</sup>R<sup>5</sup>, phenyl, -NO<sub>2</sub>, -CO<sub>2</sub>R<sup>4</sup>, -CON(R<sup>4</sup>)<sub>2</sub> wherein each R<sup>4</sup> is the same or different, or -CN;

(15) R<sup>12</sup> is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

(16) R<sup>13</sup> is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

20 (17) a (subscript for R<sup>12</sup>) is 0 to 2;

(18) b (subscript for R<sup>13</sup>) is 0 to 2;

(19) c (subscript for R<sup>6</sup>) is 0 to 2;

(20) e is 0 to 5;

(21) m is 1 or 2;

25 (22) n is 1, 2 or 3; and

(23) p is 1, 2 or 3, with the proviso that when M<sup>3</sup> and M<sup>4</sup> are both nitrogen, then p is 2 or 3 (i.e., p is not 1 when M<sup>3</sup> and M<sup>2</sup> are both nitrogen).

30 This invention also provides a pharmaceutical composition comprising an effective amount of compound of Formula I, and a pharmaceutically acceptable carrier.

This invention further provides a method of treating: allergy, allergy-induced airway (e.g., upper airway) responses, congestion (e.g., nasal congestion),

- 5 -

hypotension, cardiovascular disease, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders (e.g., hypersomnia, somnolence, and narcolepsy), disturbances of the central nervous system, attention deficit hyperactivity disorder ADHD), hypo and hyperactivity of the central nervous system (for example, agitation and depression), and other CNS disorders (such as Alzheimer's, schizophrenia, and migraine) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a method of treating: allergy comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a method of treating: allergy-induced airway (e.g., upper airway) responses comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a method of treating: congestion (e.g., nasal congestion) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a pharmaceutical composition comprising an effective amount of a compound of Formula I, and an effective amount of a H<sub>1</sub> receptor antagonist in combination with a pharmaceutically acceptable carrier.

This invention further provides a method of treating: allergy, allergy-induced airway (e.g., upper airway) responses, and congestion (e.g., nasal congestion) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

This invention further provides a method of treating: allergy comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

This invention further provides a method of treating: allergy-induced airway (e.g., upper airway) responses comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a

- 6 -

compound of Formula I in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

This invention further provides a method of treating: congestion (e.g., nasal congestion) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

#### DETAILED DESCRIPTION OF THE INVENTION

As used herein, the following terms have the following meanings, unless indicated otherwise:

alkyl-(including the alkyl portions of alkoxy and alkylaryl)-represents straight and branched carbon chains and contains from one to twenty carbon atoms, preferably one to six carbon atoms;

alkylaryl-represents an alkyl group, as defined above, bound to an aryl group, as defined below, wherein said aryl group is bound to the rest of the molecule;

aryl (including the aryl portion of alkylaryl)-represents a carbocyclic group containing from 6 to 15 carbon atoms and having at least one aromatic ring (e.g., aryl is a phenyl ring), with all available substitutable carbon atoms of the carbocyclic group being intended as possible points of attachment;

arylalkyl-represents an aryl group, as defined above, bound to an alkyl group, as defined above, wherein said alkyl group is bound to the rest of the molecule;


cycloalkyl-represents saturated carbocyclic rings of from 3 to 20 carbon atoms, preferably 3 to 7 carbon atoms;

halo (halogen)-represents fluoro, chloro, bromo and iodo;

heteroaryl-represents cyclic groups, having at least one heteroatom selected from O, S or N, said heteroatom interrupting a carbocyclic ring structure and having a sufficient number of delocalized pi electrons to provide aromatic character, with the aromatic heterocyclic groups preferably containing from 2 to 14 carbon atoms; examples include but are not limited to isothiazolyl, isoxazolyl, furazanyl, triazolyl, thiazolyl, thienyl, furanyl (furyl), pyrrolyl, pyrazolyl, pyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyridyl (e.g., 2-, 3-, or 4-pyridyl), pyridyl N-oxide (e.g., 2-, 3-, or 4-pyridyl N-oxide), triazinyl, pteridinyl, indolyl (benzopyrrolyl), pyridopyrazinyl, isoquinolinyl,

- 7 -

quinolinyl, quinoxolinyl, naphthyridinyl, wherein said pyridyl N-oxide can be represented as:




heterocycloalkyl-represents a saturated, carbocyclic ring containing from 3 to 15 carbon atoms, preferably from 4 to 6 carbon atoms, which carbocyclic ring is interrupted by 1 to 3 hetero groups selected from -O-, -S- or -NR<sup>40</sup>- wherein R<sup>40</sup> represents C<sub>1</sub> to C<sub>6</sub> alkyl, arylalkyl, -C(O)R<sup>4</sup>, -C(O)OR<sup>4</sup>, or -C(O)N(R<sup>45</sup>)<sub>2</sub> (wherein R<sup>45</sup> is as defined above, and each R<sup>45</sup> is independently selected); examples include but are not limited to 2- or 3-tetrahydrofuranyl, 2- or 3- tetrahydrothienyl, 2-, 3- or 4-piperidiny, 2- or 3-pyrrolidiny, 2- or 3-piperiziny, 2- or 4-dioxany, 1,3-dioxolany, 1,3,5-trithianyl, pentamethylene sulfide, perhydroisoquinolinyl, decahydroquinolinyl, trimethylene oxide, azetidiny, 1-azacycloheptany, 1,3-dithianyl, 1,3,5-trioxany, morpholinyl, thiomorpholinyl, 1,4-thioxany, and 1,3,5-hexahydrotriazinyl, thiazolidinyl, tetrahydropyrany;


lower alkyl-represents an alkyl group, as defined above, that comprises 1 to 6 carbon atoms, preferably 1-4 carbon atoms;

lower alkoxy-represents an alkoxy group whose alkyl moiety comprises 1 to 6 carbon atoms, preferably 1-4 carbon atoms;

=C(O)-represents



=C(NOR<sup>3</sup>)-represents




wherein (1) represents a mixture of oxime isomers; (2) represents one geometric isomer of the oxime wherein the -OR<sup>3</sup> group is on the same side of the double bond as the group to the left of the carbon atom; (3) represents one geometric isomer of the




- 8 -


oxime wherein the  $-OR^3$  group is on the same side of the double bond as the group to the right of the carbon atom; and (1) can also be represented as:




$=C(NNR^4R^5)$  represents




and represents a mixture of the isomers




$-(N)C(NR^4R^5)_2$  represents



(N) in the structure



represents a nitrogen atom that is located at one of the 4 non-fused positions of the ring, i.e., positions 1, 2, 3 or 4 indicated below:



15. AcOH-represents acetic acid;  
 t-BOC-represents t-butyloxycarbonyl;  
 Ci/mmol-represents curie/mmol (a measure of specific activity);  
 m-CPBA-represents m-chloroperbenzoic acid;  
 CSA-represents camphorsulfonic acid;  
 20 CBZ-represents carbonylbenzyloxy ( $-\text{C}(\text{O})\text{OCH}_2\text{C}_6\text{H}_5$ );

- 9 -

DBU-represents 1,8-diazabicyclo[5.4.0]undec-7-ene;  
DBN-represents 1,5-diazabicyclo[4.3.0]non-5-ene;  
DCC-represents dicyclohexylcarbodiimide;  
Dibal-H-represents diisobutylaluminum hydride;  
5 DIPEA-represents N,N-diisopropylethylamine;  
DMAP-represents 4-(dimethylamino)pyridine;  
DEC-represents 2-diethylaminoethyl chloride hydrochloride;  
DMF-represents dimethylformamide;  
EDCI-represents 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide;  
10 EtOAc-represents ethyl acetate;  
EtOH-represents ethanol;  
Fmoc-represents 9-fluorenylmethoxycarbonyl;  
HOBT-represents 1-hydroxybenzotriazole;  
HPLC-represents high performance liquid chromatography;  
15 HRMS-represents high resolution mass spectrometry;  
K<sub>i</sub>-represents inhibition constant for substrate/receptor complex;  
LAH-lithium aluminum hydride;  
LDA-represents lithium diisopropylamide;  
LRMS-represents low resolution mass spectrometry;  
20 MeOH-represents methanol;  
NaBH(OAc)<sub>3</sub>-represents sodium triacetoxyborohydride;  
NaBH<sub>4</sub>-represents sodium borohydride;  
NaBH<sub>3</sub>CN-represents sodium cyanoborohydride;  
NaHMDS-represents sodium hexamethyl disilylazide;  
25 nM-represents nanomolar;  
pA<sub>2</sub>-represents -logEC<sub>50</sub>, as defined by J. Hey, Eur. J. Pharmacol., (1995),  
Vol. 294, 329-335;  
PCC-represents pyridinium chlorochromate ;  
PyBOP-represents benzotriazole-1-yl-oxy-trispyrrolidino-phosphonium  
30 hexafluorophosphate;  
TEMPO-represents 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical ;  
TFA-represents trifluoroacetic acid;  
TMAD-represents N,N,N',N'-tetramethylazodicarboxamide;

- 10 -

TMEDA-represents tetramethylethylenediamine;  
Tr-represents triphenylmethyl;  
Tris-represents tris(hydroxymethyl)aminomethane;and  
p-TsOH-represents p-toluenesulfonic acid.

5

Also, as used herein, "upper airway" usually means the upper respiratory system—i.e., the nose, throat, and associated structures.

Also, as used herein, "effective amount" generally means a therapeutically effective amount.

10 Lines drawn into the rings indicate that the indicated bond may be attached to any of the substitutable ring carbon atoms.

Certain compounds of the invention may exist in different isomeric (e.g., enantiomers, diastereoisomers and geometric) forms. The invention contemplates all such isomers both in pure form and in admixture, including racemic mixtures. Enol  
15 forms are also included.

The compounds of this invention are ligands for the histamine H<sub>3</sub> receptor. The compounds of this invention can also be described as antagonists of the H<sub>3</sub> receptor, or as H<sub>3</sub> antagonists.

The compounds of the invention are basic and form pharmaceutically  
20 acceptable salts with organic and inorganic acids. Examples of suitable acids for such salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid  
25 to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their corresponding salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the salts are  
30 otherwise equivalent to their corresponding free base forms for purposes of this invention.

The compounds of Formula I can exist in unsolvated as well as solvated forms, including hydrated forms, e.g., hemi-hydrate. In general, the solvated forms, with

- 11 -

pharmaceutically acceptable solvents such as water, ethanol and the like are equivalent to the unsolvated forms for purposes of the invention.

The compounds of this invention can be combined with an H<sub>1</sub> receptor antagonist (i.e., the compounds of this invention can be combined with an H<sub>1</sub> receptor antagonist in a pharmaceutical composition, or the compounds of this invention can be administered with H<sub>1</sub> receptor antagonist).

Numerous chemical substances are known to have histamine H<sub>1</sub> receptor antagonist activity. Many useful compounds can be classified as ethanolamines, ethylenediamines, alkylamines, phenothiazines or piperidines. Representative H<sub>1</sub> receptor antagonists include, without limitation: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine (also known as SCH-34117), diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine and triprolidine. Other compounds can readily be evaluated to determine activity at H<sub>1</sub> receptors by known methods, including specific blockade of the contractile response to histamine of isolated guinea pig ileum. See for example, WO98/06394 published February 19, 1998.

Thus, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H<sub>1</sub> receptor antagonist, said H<sub>1</sub> receptor antagonist is selected from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or triprolidine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H<sub>1</sub> receptor antagonist, said H<sub>1</sub> receptor antagonist is selected from: astemizole, azatadine, azelastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, carebastine, descarboethoxyloratadine,

- 12 -

diphenhydramine, doxylamine, ebastine, fexofenadine, loratadine, levocabastine, mizolastine, norastemizole, or terfenadine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H<sub>1</sub> receptor antagonist, said H<sub>1</sub> receptor antagonist is selected from: azatadine, brompheniramine, cetirizine, chlorpheniramine, carebastine, descarboethoxyloratadine (also known as SCH-34117), diphenhydramine, ebastine, fexofenadine, loratadine, or norastemizole.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H<sub>1</sub> receptor antagonist, said H<sub>1</sub> receptor antagonist is loratadine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H<sub>1</sub> receptor antagonist, said H<sub>1</sub> receptor antagonist is descarboethoxyloratadine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H<sub>1</sub> receptor antagonist, said H<sub>1</sub> receptor antagonist is fexofenadine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H<sub>1</sub> receptor antagonist, said H<sub>1</sub> receptor antagonist is cetirizine.

Preferably, in the above methods, allergy-induced airway responses are treated.

Also, preferably, in the above methods, allergy is treated.

Also, preferably, in the above methods, nasal congestion is treated.

Preferably, in the above methods using a combination of a compound of Formula I (H<sub>3</sub> antagonist) and an H<sub>1</sub> antagonist, the H<sub>1</sub> antagonist is selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine. Most preferably the H<sub>1</sub> antagonist is loratadine or descarboethoxyloratadine.

In the methods of this invention wherein a combination of an H<sub>3</sub> antagonist of this invention (compound of Formula I) is administered with a H<sub>1</sub> antagonist, the antagonists can be administered simultaneously, consecutively (one after the other within a relatively short period of time), or sequentially (first one and then the other over a period of time). In general, when the antagonists are administered

- 13 -

consecutively or sequentially, the H<sub>3</sub> antagonist of this invention (compound of Formula I) is administered first.

Thus, one embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 32 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 54 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 55 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 253A and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 287 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 320 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 32.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity

- 14 -

of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 54.

Another embodiment of this invention is directed to a method of treating:

5 allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine  
10 comprising administering to a patient in need of such treatment an effective amount of Compound 55.

Another embodiment of this invention is directed to a method of treating:

allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic  
15 secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 253A.

20 Another embodiment of this invention is directed to a method of treating:

allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity  
25 of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 287.

Another embodiment of this invention is directed to a method of treating:

allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular  
30 disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine

- 15 -

comprising administering to a patient in need of such treatment an effective amount of Compound 320.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of  
5 such treatment an effective amount of Compound 32.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 54.

Another embodiment of this invention is directed to a method of treating  
10 allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 55.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 253A.

15 Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 287.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of  
20 such treatment an effective amount of Compound 320.

Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32.

Another embodiment of this invention is directed to a method of treating allergy  
25 or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54.

Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55.

30 Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A.



- 16 -

Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287.

5 Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 320.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 32, and an effective amount of H<sub>1</sub> receptor antagonist, and a pharmaceutically effective carrier.

10 Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 54, and an effective amount of H<sub>1</sub> receptor antagonist, and a pharmaceutically effective carrier. Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 55, and an effective amount of H<sub>1</sub> receptor  
15 antagonist, and a pharmaceutically effective carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 253A, and an effective amount of H<sub>1</sub> receptor antagonist, and a pharmaceutically effective carrier.

20 Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 287, and an effective amount of H<sub>1</sub> receptor antagonist, and a pharmaceutically effective carrier. Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 320, and an effective amount of H<sub>1</sub> receptor antagonist, and a pharmaceutically effective carrier.

25 Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

30 Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

- 17 -

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of a Compound 55 in combination with an effective amount of an H<sub>1</sub> receptor antagonist. Another  
5 embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

Another embodiment of this invention is directed to a method of treating:  
10 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287 in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering  
15 to a patient in need of such treatment an effective amount of Compound 320 in combination with an effective amount of an H<sub>1</sub> receptor antagonist.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in  
20 combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine,  
25 meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripeleminamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering  
30 to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine,

- 18 -

descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or  
5 triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:

10 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast,  
15 pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:

20 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast,  
25 pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:

30 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine,

- 19 -

descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripeleminamine, temelastine, trimeprazine or  
5 triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 320 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:  
10 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast,  
15 pyrilamine, promethazine, terfenadine, tripeleminamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:  
20 loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:  
25 loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:  
30 loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering

- 20 -

to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating:  
5 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating:  
10 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 320 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating:  
15 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:  
20 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:  
25 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55 in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:  
30 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from: loratadine or descarboethoxyloratadine.

- 21 -


Another embodiment of this invention is directed to a method of treating:  
allergy, allergy-induced airway responses, and congestion comprising administering  
to a patient in need of such treatment an effective amount of Compound 287 in  
combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:  
5 loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:  
allergy, allergy-induced airway responses, and congestion comprising administering  
to a patient in need of such treatment an effective amount of Compound 320 in  
combination with an effective amount of an H<sub>1</sub> receptor antagonist selected from:  
10 loratadine or descarboethoxyloratadine.


R<sup>1</sup> is preferably selected from:

- (A) aryl (most preferably phenyl);
- (B) substituted aryl (e.g., substituted phenyl), wherein the  
15 substituents on said substituted aryl are most preferably selected from: (1) halo (e.g.,  
monohalo or dihalo), more preferably chloro or fluoro, even more preferably  
monochloro, dichloro, monofluoro or difluoro; or (2) alkyl, more preferably unbranched  
(i.e., straight chain, e.g., methyl) alkyl, even more preferably substituted alkyl, still  
more preferably alkyl substituted with halo (e.g., 1, 2 or 3 halo atoms, such as Cl or  
20 F), even still more preferably alkyl substituted with fluoro atoms, yet still more  
preferably trifluoromethyl;

- (C) heteroaryl, most preferably a five or six membered heteroaryl  
ring, more preferably a six membered heteroaryl ring, and still more preferably pyridyl,  
examples of heteroaryl rings include pyridyl, thienyl, pyrimidinyl, thiazolyl or pyridyl N-  
25 Oxide, most preferred heteroaryl rings are exemplified by




wherein




- 22 -

is preferred more;


- (D) substituted heteroaryl, most preferably halo or alkyl substituted heteroaryl (e.g., halopyridyl (e.g., fluoropyridyl) and alkylthiazolyl), more preferably substituted heteroaryl wherein the substituents are independently selected from the same or different alkyl groups (even more preferably one straight chain alkyl group, e.g., methyl), still more preferably alkyl substituted thiazolyl, and even more preferably



yet even more preferably



- 10 (E) when R<sup>1</sup> is taken together with X, then the moiety is



wherein c is most preferably 0 or 1, and when c is 1 then R<sup>6</sup> is most preferably halo, and when c is 1 then R<sup>6</sup> is more preferably fluoro.

- X is preferably =C(NOR<sup>3</sup>) wherein R<sup>3</sup> is preferably selected from H, alkyl or halo substituted alkyl (e.g., fluoro substituted alkyl, such as -CH<sub>2</sub>CF<sub>3</sub>), most preferably alkyl, more preferably methyl or ethyl, and still more preferably methyl.

Preferably M<sup>2</sup> is nitrogen.

n is preferably 2.

a is preferably 0 or 1, and most preferably 0.

- 20 b is preferably 0 or 1, and most preferably 0.

c is preferably 0 or 1, and most preferably 0, and when c is 1 then R<sup>6</sup> is preferably halo, and when c is 1 R<sup>6</sup> is most preferably fluoro.

e is preferably 1-5.


Y is preferably =C(O) (i.e., =C=O).

- 25 M<sup>3</sup> and M<sup>4</sup> are preferably selected such that: (1) one is carbon and the other is nitrogen, or (2) both are nitrogen, with M<sup>3</sup> most preferably being carbon.


- 23 -

p is preferably 2.


Z is preferably C<sub>1</sub> to C<sub>3</sub> alkyl, and most preferably



R<sup>2</sup> is preferably a six membered heteroaryl ring, most preferably pyridyl, substituted pyridyl, pyrimidinyl or substituted pyrimidinyl, more preferably pyridyl, pyridyl substituted with —NR<sup>4</sup>R<sup>5</sup>, pyrimidinyl or pyrimidinyl substituted with —NR<sup>4</sup>R<sup>5</sup>, still more preferably pyridyl, pyridyl substituted with —NH<sub>2</sub> (i.e., R<sup>4</sup> and R<sup>5</sup> are H), pyrimidinyl or pyrimidinyl substituted with —NH<sub>2</sub> (i.e., R<sup>4</sup> and R<sup>5</sup> are H), and even more preferably



and still even more preferably



R<sup>3</sup> is preferably H or alkyl, most preferably H or methyl.

R<sup>4</sup> is preferably H or lower alkyl, most preferably H or methyl, and more preferably H.

R<sup>5</sup> is preferably H, C<sub>1</sub> to C<sub>6</sub>alkyl or —C(O)R<sup>4</sup>, most preferably H or methyl, and more preferably H.

R<sup>12</sup> is preferably alkyl, hydroxyl or fluoro, and most preferably H.

R<sup>13</sup> is preferably alkyl, hydroxyl or fluoro, and most preferably H.

Representative compounds of this invention include, but are not limited to: Compounds 23, 30, 31, 32, 33, 41, 44, 45, 49, 50, 52, 53, 54, 55, 56, 57A, 59, 65, 75, 76, 80, 82, 83, 88, 92, 99, 104, 105, 110, 111, 117, 121, 123, 127, 128, 200-241, 244-273, 275, and 278-282, 287, 296, 301-439 and 446.

Thus, representative compounds of this invention include, but are not limited to: Compounds 23, 30, 31, 32, 33, 44, 45, 49, 50, 53, 54, 55, 59, 75, 76, 83, 88, 92,



- 24 -

99, 104, 110, 117, 128, 200, 201, 203-215, 217-241, 244-246, 246A, 247-253, 253A, 254-273, 275, 278, and 280-282, 317, 334 and 403.

Preferred compounds of this invention are selected from: Compound 23, 30, 31, 32, 33, 50, 53, 54, 55, 56, 57A, 59, 92, 212, 215, 218, 219, 220, 224, 225, 226, 227, 229, 233, 235, 237, 238, 246, 246A, 247, 248, 251, 253, 253A, 268-273, 275, 278-281, 287, 296, 301, 304-307, 309, 312, 314-318, 320-356, or 358-376.

Most preferred compounds of this invention are selected from: Compound 30, 31, 32, 33, 54, 55, 56, 57A, 225, 237, 246A, 253A, 273, 280, 287, 296, 301, 304-307, 309, 312, 314-318, 320-348, 350-356, 359-372, and 374-376.

Thus, one embodiment of this invention is directed to Compound 32.

Another embodiment of this invention is directed to Compound 54.

Another embodiment of this invention is directed to Compound 55.


Another embodiment of this invention is directed to Compound 253A.

Another embodiment of this invention is directed to Compound 287.


Another embodiment of this invention is directed to Compound 320.

Structures for the above compounds are found in the Examples below, and in Tables 1 to 3 below.

The more preferred compound of this invention is the compound of the formula:




This invention also provides a compound of the formula:



This invention also provides a compound of the formula:


- 25 -



Compounds 32A and 32B can also be used in the pharmaceutical compositions, and the methods of this invention.


The following processes may be employed to produce compounds of the invention.

One synthetic route involves a linear sequence of reactions to obtain the desired compounds, i.e.,




This linear sequence of reactions to synthesize compounds of this invention is illustrated below. In the illustrated procedure  $R^1$  is aryl, heteroaryl, or alkyl;  $X$  = a ketone, oxime or substituted oxime;  $M^1 = M^3$  = carbon;  $M^2 = M^4$  = nitrogen;  $Y$  is  $C=O$ ;  $Z = CHR$ ;  $R^2$  is heteroaryl; and  $n$  and  $m = 2$  ( $n$  and  $m$  being 1 can also be prepared by this procedure).

- 26 -

**Step: 1 Synthesis of Ketone 8**

In the above equations PG represents a protecting group, and M represents Li or  $MgX^1$  (wherein  $X^1$  represents Cl, Br or I).


In equation 1 and 2, a Grignard reagent 2 is reacted with an electrophile such as the aldehyde 1 or the nitrile 4 in a suitable aprotic solvent such as THF or ether. PG represents a protecting group. Suitable protecting groups include, for example, methyl and benzyl. In the case of nitrile 4, acidic workup yields the ketone 8 directly. Alcohol 3 can be oxidized by a number of different reagents to give 8. Alternatively, the amide 7 can be reacted with an organometallic reagent to directly give the ketone 8. Suitable protecting groups for this step include carbamates or amides or the like. Thus, examples of protecting groups in equation 3 include t-BOC, CBZ and Fmoc.

**Step 2: Deprotection of 8**

- 27 -


When the protecting group, PG, is a methyl group, said methyl group can be removed using a reagent such as a chloroformate; when PG is a carbamate, such as, a t-Boc group, it can be removed by dilute acid, such as, for example HCl.

### 5 Step 3: Synthesis of 11




Amine 9 can be coupled to acid 10 using a number of methods well known in the art such as DCC or PyBOP. Alternatively, the acid 10 can be activated by conversion to the acid chloride or mixed anhydride and then reacted with the amine 9 to give 11. Suitable protecting groups for 10 include, for example, t-Boc.

### Step 4: Synthesis of Amine 12



Compound 11 in which the protecting group is a t-Boc can be deprotected under acidic conditions such as HCl in dioxane or TFA in CH<sub>2</sub>Cl<sub>2</sub> to give the amine 12.

### Step 5: Synthesis of Compound 14




R<sup>30</sup> in 13 represents an alkyl group. E is a leaving group, halogen, or E is a carbonyl group.

Compound 14 can be prepared by reacting amine 12 with 13. When E represents a carbonyl group (C=O), 12 and 13 are combined in a solvent such as

- 28 -




CH<sub>2</sub>Cl<sub>2</sub> in the presence of molecular sieves. After the reaction is complete (e.g., 1 to 10 h), a reducing agent such as NaBH(OAc)<sub>3</sub> is added. Alternatively, when E is a halogen atom such as Cl or Br, 12 and 13 are combined in a solvent, such as DMF, in the presence of a tertiary amine base to give the product 14. Suitable protecting groups include, for example t-Boc, phthaloyl.

#### Step 6: Synthesis of Compound 16




Compound 14 can be converted to the oxime 15 by combining 14 with H<sub>2</sub>NOR<sup>3</sup>·HCl in pyridine at a temperature of 40 – 60°C. Alternatively, 14 can be combined with H<sub>2</sub>NOR<sup>3</sup>·HCl in an alcoholic solvent in the presence of a base, such as, NaOAc, to give 15.

An alternate approach to the synthesis of compounds of Formula I involves the synthesis of the two halves of the molecule followed by coupling of the two pieces, i.e.,



In this case, the synthesis of the AB fragment is the same as that described above. The synthesis of the CD fragment is given below.

#### Step 1: Synthesis of Compound 17




R<sup>30</sup> is as defined above (i.e., alkyl). R<sup>35</sup> is methyl or ethyl.

- 29 -

Compound 17 is synthesized in the same manner as that described for the synthesis of compound 14.

### Step 2: Synthesis of Compound 18



5

M represents Li, Na, or K.


Compound 17 is saponified in a mixed solvent, such as, for example: (1) EtOH or MeOH and water, or (2) THF, water, and MeOH, using an alkali metal base such as LiOH or NaOH at a temperature of from 50 to 100°C to give the salt 18.

Compound 18 can be combined with compound 9, as described above, to give 14. The remaining steps are the same.

Compounds useful in this invention are exemplified by the following examples which should not be construed as limiting the scope of the disclosure. Alternative mechanistic pathways and analogous structures within the scope of the invention may be apparent to those skilled in the art.

### Example 1

#### Step 1




1A


To a solution of 10.81 g (100 mmol) of 2-amino-4-methylpyridine in 250 ml of tert-butanol was added 26.19 g (120 mmol) of BOC anhydride. The reaction mixture was stirred at room temperature overnight, concentrated, loaded on silica gel and flash chromatographed (from 30% hexanes/ CH<sub>2</sub>Cl<sub>2</sub> to 0 - 2% acetone/ CH<sub>2</sub>Cl<sub>2</sub>) to produce 15.25 g (73.32 mmol; 73%) of 1A as a white solid.

25

- 30 -

**Step 2**

To a  $-78^{\circ}\text{C}$  solution of 1A (35.96 g, 173 mmol) in THF (1.4 L) was added 1.4 M BuLi solution (272 ml, 381 mmol) in hexanes in portions over 30 min. Reaction mixture was then allowed to warm up and was stirred for 2 h at room temperature, which resulted in the formation of an orange precipitate. The mixture was cooled back to  $-78^{\circ}\text{C}$ , and predried oxygen (passed through a Drierite column) was bubbled through the suspension for 6 h while the temperature was maintained at  $-78^{\circ}\text{C}$ . Reaction mixture color changed to yellow during this time. It was then quenched at  $-78^{\circ}\text{C}$  with 51.4 ml (700 mmol) of  $\text{Me}_2\text{S}$  followed by 22 ml (384 mmol) of AcOH. Reaction mixture was allowed to warm up and was stirred for 48 h at room temperature. Dilution with water and extraction with EtOAc were followed by concentration and flash chromatography (0 – 15% acetone/  $\text{CH}_2\text{Cl}_2$ ) to provide 20.15 g (90 mmol; 52%) of alcohol 2A as a pale yellow solid.


**Step 3**

To a solution of 19.15 g (85.5 mmol) of alcohol 2A in 640 ml of  $\text{CH}_2\text{Cl}_2$  was added saturated aqueous solution of 8.62 g (103 mmol) of  $\text{NaHCO}_3$  and 444 mg (4.3 mmol) of NaBr. Reaction mixture was cooled to  $0^{\circ}\text{C}$ , and 140 mg (0.90 mmol) of TEMPO was introduced. Upon vigorous stirring 122 ml of 0.7 M (85.4 mmol) commercial bleach solution (5.25% in NaOCl) was added in portions over 40 min. After additional 20 min at  $0^{\circ}\text{C}$  reaction mixture was quenched with saturated aqueous  $\text{Na}_2\text{S}_2\text{O}_3$  and allowed to warm to room temperature. Dilution with water and

- 31 -


extraction with  $\text{CH}_2\text{Cl}_2$  were followed by concentration and flash chromatography (from 30% hexanes/  $\text{CH}_2\text{Cl}_2$  to 0 - 2% acetone/  $\text{CH}_2\text{Cl}_2$ ) to afford 15.97 g (71.9 mmol; 84%) of aldehyde **3A** as an off-white solid.

5 **Step 4**



To a solution of 11.87 g (53.5 mmol) of aldehyde **3A** in 370 ml of  $\text{CH}_2\text{Cl}_2$  was added 9.07 ml (58.8 mmol) of ethyl isonipecotat followed by four drops of AcOH. Reaction mixture was then stirred for 40 min at room temperature after which 22.68 g (107 mmol) of  $\text{NaBH}(\text{OAc})_3$  was introduced. Reaction mixture was stirred overnight at room temperature, neutralized with saturated aqueous  $\text{NaHCO}_3$ , diluted with water and extracted with  $\text{CH}_2\text{Cl}_2$ . Concentration and flash chromatography (0 – 4% sat.  $\text{NH}_3$  in  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) provided 19.09 mg (52.6 mmol; 98%) of **4A** as an off-white solid.


15 **Step 5**




To a solution of 1.57 g (4.33 mmol) of ester **4A** in 10 ml of a 3 : 1 : 1 mixture of THF – water – methanol was added 0.125 g (5.21 mmol) of LiOH. Reaction mixture was stirred overnight at room temperature, concentrated and exposed to high vacuum to obtain 1.59 g of crude acid **5A** as a yellowish solid which was used without purification.



- 32 -


**Example 2**

A solution of compound **6A** (42mmol), NBS (126mmol) and Bz<sub>2</sub>O<sub>2</sub> (4.2mmol) in CCl<sub>4</sub> (400ml) was refluxed at 80°C for 5 h, cooled and stirred at room temperature overnight. The reaction was filtered and concentrated, and the residue was purified by flash column (30% EtOAc/Hexane) to obtain the desired compound **7A** (3.1g, 23%).

**Example 3****Step 1**


To a solution of **8A** (10 g, 79.4 mmol) and DMAP (0.029 g, 0.24 mmol) in methylene chloride (150 mL) at 0°C was added phthaloyl dichloride (16.1 g, 79.4 mmol) dropwise. The reaction mixture was stirred at room temperature overnight. After stirring overnight, the reaction was washed with saturated aqueous NaHCO<sub>3</sub>, water, dried and concentrated to give compound **9A** as a yellow solid (20 g, 99.8%) which was used without further purification.

- 33 -

**Step 2**


In a manner similar to that described in Example 2, compound **9A** (20 g, 79.3 mmol) was converted to compound **10A**.

5

**Step 3**

Compound **10A** (0.5 g, 1.5 mmol) and hydrazine (0.5 M in ethanol, 5 mL, 2.5 mmol) were combined and stirred at room temperature overnight. The reaction was diluted with water and extracted with methylene chloride. The organic layer was dried, concentrated and the residue purified on a flash column (3% methanol in ethyl acetate) to give compound **11A** (0.2 g, 66%).

10


**Example 4****Step 1**

15

- 34 -

Compounds **12A** (2 g, 18.3 mmol) and **13A** (3.5 g, 22 mmol) were dissolved in methylene chloride and stirred at room temperature for 1 h. NaB(OAc)<sub>3</sub>H (5.4 g, 25.6 mmol) was added and the mixture stirred at room temperature for 5h. The reaction was washed with saturated aqueous NaHCO<sub>3</sub>, dried and concentrated, and the residue purified by flash column (2% methanol in ethyl acetate). Compound **14A** was obtained (4.5g, 99%).


### Step 2



In a manner similar to that described in Example 1, Step 5, compound **14A** (0.35 g, 1.4 mmol) was converted to compound **15A** (0.31g, 100%).


### Example 5

#### Step 1




To the solution of 2,4-diflorobenzaldehyde (**16A**, 28.1mmol) in THF (10ml) was added the Grignard reagent **17A** ( 1.33M in THF, 30ml), and the mixture was stirred at room temperature overnight. The reaction was quenched with saturated NH<sub>4</sub>Cl (150ml), extracted three times with EtOAc (100ml), dried, filtered and concentrated. Flash chromatography (20% MeOH/EtOAc) yielded the desired compound **18A** (1.8g, 27%).


- 35 -

**Step 2**

Compound **18A** (1.6 g, 6.7 mmol),  $\text{H}_2\text{NHOH}\cdot\text{HCl}$  (0.95 g, 6.7 mmol) and pyridine (10 mL) were combined and heated to 60°C overnight. The pyridine was removed under vacuum and the residue treated with methylene chloride and saturated aqueous  $\text{NaHCO}_3$ . The organic layer was separated, dried, and concentrated, and the residue purified by flash chromatography to give compound **19A** (1.4g, 82%).


**Step 3**

To the suspension of NaH (0.41g, 10.2 mmol) in THF (10ml) was slowly added a solution of **19A** (1.3 g, 5.11 mmol) in DMF (5 mL) dropwise and the reaction stirred at 70~75°C overnight. The mixture was extracted twice with EtOAc and three times with  $\text{H}_2\text{O}$  (30 mL), dried over  $\text{MgSO}_4$  and concentrated to give crude **20A** which was used without further purification (1.04 g, 87%).


**Step 4**

- 36 -

To the solution of compound **20A** (4.3 mmol) in dichloroethane (20 ml) at 0°C was added 2-chloroethyl chloroformate (6.2 mmol) and triethylamine (7.2 mmol) and the reaction was stirred at room temperature overnight. The solvent was evaporated, Et<sub>2</sub>O was added to the residue, and the unreacted starting material was removed by  
5 filtration. The filtrate was concentrated and the residue redissolved in MeOH and refluxed for 30 min. Removal of the methanol gave the product **21** (0.3g) which was used without further purification.

**Step 5**

To a mixture of compound **21** (1.64 mmol), compound **5A** (1.64 mmol) and PyBOP (1.64 mmol) was added DIPEA (4.92 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (10 ml), and the reaction was stirred over the weekend at room temperature. Saturated NaHCO<sub>3</sub> (100 ml) was added and the reaction was extracted and twice with CH<sub>2</sub>Cl<sub>2</sub> (100 mL), dried  
15 over solid MgSO<sub>4</sub>, concentrated and flash chromatographed (70% EtOAc/Hexane) to give compound **22** (1.04 mmol, 64%).


**Step 6**

Compound **22** (0.2 g, 0.37 mmol) was dissolved in CF<sub>3</sub>CO<sub>2</sub>H (3 mL) and methylene chloride (3 mL) and stirred at room temperature overnight. The solvent was removed by evaporation, saturated aqueous NaHCO<sub>3</sub> was added and mixture  
20

extracted with methylene chloride. The organic layer was dried ( $\text{MgSO}_4$ ), filtered and concentrated, and the residue purified by flash chromatography to give compound **23** (0.11 g, 68%).


### 5 Example 6

### Step 1




A solution of **24** (50 g, 387 mmol) and triethylamine (110 mL) in dioxane (400 mL) and water (400 mL) at 4°C was treated with Boc<sub>2</sub>O (93 g, 426 mmol). The cooling bath was removed and the solution allowed to warm to room temperature. After 21h, the volume was reduced by two-thirds under vacuum. The residue was poured into ethyl acetate (250 mL) and water (250 mL). Saturated aqueous NaHCO<sub>3</sub> (250 mL) was added and the organic phase was separated and discarded. The aqueous phase was acidified with 10% HCl and extracted with ethyl acetate. The combined organic phases were washed with water, brine, and dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to give **25** as a white powder (82 g, 94%).

### Step 2




To a solution of compound **25** (40 g, 175 mmol) in DMF (250 mL) at 4°C was added N,O – dimethylhydroxylamine, hydrochloride (34 g), EDCI (44 g, 0.228 mol), HOBT (2.4 g), and DIPEA (120 mL). The reaction was warmed to room temperature and stirred overnight. The reaction was then concentrated to half volume in vacuo and poured onto 1:1 ethyl acetate:water. The organic layer was separated and the aqueous layer extracted with additional ethyl acetate. The combined organic layers were washed with saturated aqueous NH<sub>4</sub>Cl, saturated aqueous NaHCO<sub>3</sub>, water, and brine, and dried. Concentration gave **26** as a light yellow oil (46.7 g, 99%)

- 38 -


**Step 3**

To a solution of 2-bromopyridine (17.6 mL, 0.184 mol) in THF (600 mL) at  $-78^{\circ}\text{C}$  was added n-BuLi (115 mL of a 1.6M solution in hexanes, 0.184 mol) dropwise over 15 min. After stirring for an additional 30 min at this temperature, a solution of **26** (25 g, 91.9 mmol) in THF (500 mL) was added dropwise over 15 min. The reaction was removed from the cold bath and placed in an oil bath and heated to  $60^{\circ}\text{C}$  for 1.5h. The reaction was then cooled to  $4^{\circ}\text{C}$ , diluted with ether (500 mL), and treated with saturated aqueous  $\text{Na}_2\text{SO}_4$  ( $\approx 5$  mL). The mixture was transferred to an Erlenmeyer flask and diluted with additional ether (700 mL). Additional saturated aqueous  $\text{Na}_2\text{SO}_4$  was added followed by solid  $\text{Na}_2\text{SO}_4$ . The mixture was filtered through a plug of solid  $\text{Na}_2\text{SO}_4$  and concentrated in vacuo. Flash column chromatography (0-20% ethyl acetate in hexanes) yielded compound **27** as a yellow oil (16.85 g, 63%).


**Step 4**

A solution of **27** (3.3 g, 11.4 mmol) in methanol (50 mL) was treated with 4M HCl in dioxane (50 mL) and stirred at room temperature for 1.5 h. Removal of the solvent in vacuo gave **28** as a tan powder (3g, 100%).

- 39 -

**Step 5**

To a suspension of compound 5A (17.4 g, 50 mmol), compound 28 (11 g, 42 mmol), and diisopropylethylamine (34.6 mL, 199 mmol) in DMF (125 mL) was added  
5 HOBT (7.83 g, 58 mmol), EDC (18.54 g, 96.7 mmol), and 4Å molecular sieves. The mixture was stirred for 40 h at room temperature, diluted with methylene chloride (600 mL) and 0.5 N NaOH (400 mL) and filtered. The precipitate was washed thoroughly with additional 0.5N NaOH and methylene chloride. The combined organic phases were concentrated and chromatographed twice on silica gel (1:1 hexane:methylene  
10 chloride to 6% saturated NH<sub>3</sub> in methanol in methylene chloride) to produce 29 as a tan solid (22.3 g) which was used as is in the next step.


**Step 6**

15 A solution of 29 (22.3 g, 44 mmol) in methylene chloride (120 mL) and trifluoroacetic acid (60 mL) was stirred for 7 h at room temperature. The reaction was




- 40 -

concentrated, exposed to high vacuum for 3h, dissolved in toluene and concentrated and then exposed again to high vacuum. The so-obtained crude brown oil was used in the next step without further purification.

5 **Step 7**


Compound 30 ( $\approx 17.9$  g, 44 mmol) was dissolved in pyridine (420 mL), treated with  $\text{H}_2\text{NOCH}_3\cdot\text{HCl}$  (21.78 g, 264 mmol) and heated to  $90^\circ\text{C}$  for 14h. The reaction was then concentrated and the residue taken up in a mixture of methylene chloride, (500 mL) and 2N NaOH (500 mL). The organic phase was separated and the aqueous phase extracted with additional methylene chloride (300 mL). The organic phases were dried and concentrated, and the residue chromatographed on  $\text{SiO}_2$  (0-13%  $\text{NH}_3/\text{MeOH}$  in  $\text{CH}_2\text{Cl}_2$ ) to produce a yellow solid (9.26 g). The mixed fractions from the column were rechromatographed to give an additional 3.23g of the desired material. Total yield 12.49 g (65% yield over the last two steps).

- 41 -

**Step 8**


Compound 31 (1 g) in ethanol (15 mL) was separated into the pure isomers using a Chiralcel AD column (20 mm x 500 mm) (eluent: 75:25 hexane: isopropanol plus 0.5% N,N-diethylamine; flow rate: 50 mL/min; UV detection at 254 nm) to give compound 32 (0.6 g) and compound 33 (0.4 g).  $[M+H]^+$  437 for 32 and 33.

Alternatively, compound 32 can preferably be prepared from compound 5A in a manner similar to that described for compound 287 in Step 3 of Example 28.


**Example 7****Step 1**

To a solution of 34 (2.4 g, 13.5 mmol) in THF (15 mL) was added compound 35 (26 mL of a 1.3M solution) and the reaction stirred overnight at room temperature. 2N HCl was then added till the pH < 2 and the THF was removed under reduced pressure. The pH was neutralized by the addition of 1N NaOH and the aqueous phase extracted with 5% MeOH in EtOAc. The organic phase was dried, concentrated, and the residue chromatographed (20% MeOH in EtOAc) to give 36 (1.03 g, 28%).


- 42 -

**Step 2**

To a solution of **36** (1.03 g, 3.78 mmol) in 1,2-dichloroethane (30 mL) was added 1-chloroethylchloro formate (0.76 mL, 7.6 mmol) and the reaction stirred at room temperature overnight. The solvent was removed in vacuo and the residue washed with ether. Solid residue was removed by filtration and the ether removed by evaporation to give an oil which was dissolved in MeOH (15 mL) and heated to reflux for 2h. Removal of the solvent gave **37** which was used in the next step without further purification (1.4 g).


**Step 3**

Compound **37** (0.98 g, 3.78 mmol), N-Boc isonipocotic acid (0.87 g, 3.78 mmol), DEC (1.11 g, 5.7 mmol), HOBT (0.68g, 4.91 mmol) and DIPEA (3 mL) were combined in  $\text{CH}_2\text{Cl}_2$  (40 mL) and stirred overnight at room temperature. The reaction was then diluted with  $\text{CH}_2\text{Cl}_2$  and washed with saturated aqueous  $\text{NaHCO}_3$ . The organic layer was dried, concentrated and the residue chromatographed (10% hexane in EtOAc) to give **38** (1.61 g, 91%).

**Step 4**


- 43 -

Compound **38** (1.61 g, 3.43 mmol) in  $\text{CH}_2\text{Cl}_2$  (15 mL) was treated with 1N HCl in dioxane (5.2 mL) and stirred overnight at room temperature. The solvent was removed in vacuo to give **39** (1.65 g) which was used without further purification.


5 **Step 5**

Compound **39** (1.65 g, 4.01 mmol), **7** (1.29 g, 4.07 mmol) and  $\text{Et}_3\text{N}$  (1.7 mL) were combined in DMF (40 mL) and stirred at room temperature overnight. The reaction was dissolved in EtOAc and washed 4 times with water. The organic layer  
10 was dried and concentrated, and the residue purified by chromatography (5% MeOH in EtOAc) to give **40** (0.6 g, 47%).

- 44 -


**Step 6**

A solution of 40 (0.31 g, 0.51 mmol) in pyridine (5 mL) was treated with H<sub>2</sub>NOMe·HCl (0.092 g, 1.08 mmol) and heated to 60°C overnight. The reaction was  
5 diluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>, washed with saturated aqueous NaHCO<sub>3</sub>, dried, and concentrated, and the residue purified by chromatography (10-15% MeOH in EtOAc) to give 41 (0.09 g).


**Example 8****Step 1**

In a manner similar to that described in Example 7, Steps 3-4, compound 42  
was converted to compound 43.


- 45 -

**Step 2**

To a solution of **43** (2.3 g, 6.3 mmol) in  $\text{CH}_2\text{Cl}_2$  (60 mL) was added 4Å molecular sieves and 4-formylpyridine (0.68 mL, 6.9 mmol) and the mixture stirred for 3 h at room temperature.  $\text{Na}(\text{OAc})_3\text{BH}$  (2.7 g, 12.7 mmol) was then added and the reaction stirred for 1h. The reaction was quenched by the addition of  $\text{NH}_4\text{Cl}$  followed by the addition of saturated aqueous  $\text{NaHCO}_3$ . The reaction mixture was then extracted with EtOAc, and the combined organic layers were dried and concentrated to give a residue which was chromatographed (20% MeOH in EtOAc). Compound **44** was obtained (2.3 g, 87%).


**Step 3**

In a manner similar to that described in Example 7, Step 6, compound **44** was converted to compound **45**.


**Example 9****Step 1**

- 46 -

In a manner similar to that described in Example 8, Step 2, compound 46 (1.13 g, 6 mmol) was converted to compound 47 (1.7 g, 100%).


**Step 2**

In a manner similar to that described in Example 7, Step 4, compound 47 (1.7 g, 6.13 mmol) was converted to compound 48 (1.9 g, 100%).

**Step 3**


A mixture of compound 48 (0.57 g, 2 mmol) and compound 42 (0.52 g, 2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was added Et<sub>3</sub>N (1.95 mL) and the reaction cooled to -40°C. Triphosgene (0.2 g) was added and the reaction stirred at -40°C for 2 h and room temperature for 48 h. The reaction was then washed with 1N NaOH, brine, and the organic layer dried. Concentration gave a residue that was purified by column chromatography (10% MeOH in EtOAc) to give 49 (0.14 g, 55%).

15


**Step 4**

- 47 -


In a manner similar to that described in Example 7, Step 6, compound 49 (0.09 g, 0.21 mmol) was converted to compound 50.

**Example 10****Step 1**

In a manner similar to that described in Example 7, Steps 3-4, compound 28 (2.6 g, 9.9 mmol) was converted to compound 51 (1.1 g).

**Step 2**





In a manner similar to that described in Example 7, Step 5, compound 51 (1.1 g, 2.94 mmol) was reacted with compound 11 (0.59 g, 2.94 mmol) to give compound 52 (0.53 g).

**Step 3**

In a manner similar to that described in Example 6, Step 7, compound 52 (0.53 g, 1.26 mmol) was converted to compound 53 (0.48 g).




- 48 -

**Step 4**


10 In a manner similar to that described in Example 6, Step 8, the 4 diastereomers of compound **53** could be obtained using a Chiralcel AD column (75:25 hexane:EtOAc plus 0.5% Et<sub>2</sub>NH). The two faster eluting compounds (**54** and **55**) were the E-oxime isomers and the slower eluting compounds (**56** and **57A**) were the Z-oxime isomers.

- 49 -

Isomer A	54	0.12 g
Isomer B	55	0.11 g
Isomer C	56	0.08 g
Isomer D	57A	0.06 g


**Example 11****Step 1**

A solution of *n*-BuLi (4.2 mL of a 1.6 M solution in hexane) in THF (25 mL) was treated at  $-25^{\circ}\text{C}$  with  $(i\text{-Pr})_2\text{NH}$  (0.69 g, 6.8 mmol). The reaction was stirred for 1 h at  $0^{\circ}\text{C}$  and then cooled to  $-70^{\circ}\text{C}$ . Compound 4A (0.82 g, 2.26 mmol) in THF (5 mL) was added dropwise and the reaction stirred at  $-70^{\circ}\text{C}$  for 2 h and  $-50^{\circ}\text{C}$  for 2 h. The reaction was recooled to  $-70^{\circ}\text{C}$  and (1*S*)-(+)-(10-camphorsulfonyl)oxaziridine (1.04 g, 4.52 mmol) in THF (5 mL) was added. The reaction was stirred at  $-70^{\circ}\text{C}$  for 2 h and slowly warmed to room temperature overnight. The reaction was quenched by the addition of saturated aqueous  $\text{NH}_4\text{Cl}$  and extracted with EtOAc. The organic layer was dried and concentrated, and the residue purified by column chromatography (1:1 hexane:EtOAc) to give 57 (0.44 g, 51%).

**Step 2**


In a manner similar to that described in Example 1, Step 5, compound 57 (0.42 g, 1.1 mmol) was converted to compound 58 (0.4 g).

- 50 -


**Step 3**

In a manner similar to that described in Example 6, Steps 5 - 8, compound 58 (0.25 g, 0.7 mmol) was converted to compound 59 (0.1 g).

5

**Example 12****Step 1**

A solution of compound 60 (10 g, 50.7 mmol) in ether (150 mL) at  $-78^{\circ}\text{C}$  was treated sequentially with TMEDA (11.8 g, 101.4 mmol) and *s*-BuLi (58.5 mL of a 1.3M solution in hexanes, 76 mmol) and the reaction stirred at this temperature for 6 h. Neat  $\text{CH}_3\text{SO}_4\text{CH}_3$  (12.8 g, 101.4 mmol) was then added and the reaction allowed to slowly warm to room temperature overnight. Saturated aqueous NaCl was added and the organic layer was separated. The aqueous layer was extracted three times with ether and the combined organic layers were dried, concentrated, and the residue chromatographed (5% EtOAc in hexane) to give 61 (8.0 g, 75%).


**Step 2**

A solution of 61 (8 g, 37.9 mmol) in THF (40 mL) at  $0^{\circ}\text{C}$  was treated dropwise with a solution of  $\text{BH}_3 \cdot \text{THF}$  (45.4 mL of a 1.0M solution in THF, 45.4 mmol) and the


20

- 51 -

reaction allowed to slowly warm to room temperature overnight. The reaction was recooled to 0°C, EtOH (13 mL), pH = 7 buffer (25 mL) and H<sub>2</sub>O<sub>2</sub> (25 mL) was added, and the reaction allowed to stir at room temperature overnight. The solvent was then removed in vacuo and the residue poured into water and CH<sub>2</sub>Cl<sub>2</sub>. 10% aqueous NaOH (10 mL) was added and the organic layer separated. The aqueous layer was extracted with additional CH<sub>2</sub>Cl<sub>2</sub> and the combined organic layers were dried and concentrated. The residue was chromatographed (40% EtOAc in hexane) to give 62 (3 g).

10 **Step 3**


A solution of 62 (2.8 g, 12.2 mmol) in EtOAc (30 mL) and NaBr (1.26 g, 0.12 mmol) in saturated aqueous NaHCO<sub>3</sub> (30 mL) was cooled to 0°C and treated with TEMPO (0.02 g, 0.12 mmol). After 15 min., NaOCl (17.44 mL) was added and the mixture stirred for 3 h. Saturated aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> was added and the pH adjusted to 5 – 6 by the addition of 1N HCl. The mixture was extracted with EtOAc and the organic layers were dried and concentrated. The residue was chromatographed (10 – 20% EtOAc in hexane) to give compound 63 (2.1 g, 76%).

20 **Step 4**


To a cooled (0°C) suspension of PCC (0.95 g, 4.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added dropwise a solution of 63 (0.5 g, 2.2 mmol). And the mixture stirred overnight at room temperature. Additional PCC (1 eq.) was added and the mixture was heated

- 52 -

to reflux for 2 h. The reaction was cooled, filtered through celite, and concentrated to give crude **64** (1.5 g) which was used without further purification.


**Step 5**

In a manner similar to that described in Example 5, Step 5, Example 7, Step 4, Example 1, Step 4, and Example 6, Steps 6 and 7, **64** (0.73 g, 3 mmol) was converted to **65** (0.1 g).


**Example 13****Step 1**

To a  $0^\circ\text{C}$  solution of Vilsmeier salt, prepared by the dropwise addition of phosphorus oxychloride (150.0 mL; 1.61 mol) to DMF (310.4 mL; 4.01 mol) over 15 min. and subsequent cooling in an ice bath, was added malonic acid (40.1 g; 0.39 mol) in portions over 45 min. The reaction mixture was then heated to  $100^\circ\text{C}$ , and the stirring was continued for 48 h. The reaction mixture was then allowed to cool to room temperature and was quenched by slowly pouring it into a suspension of  $\text{NaHCO}_3$  (808 g; 9.62 mol) in water. The solution was decanted off the excess of  $\text{NaHCO}_3$  and concentrated to dryness under vacuum. After exposure to high vacuum for 2 days, the solid residue was washed repeatedly with  $\text{CH}_2\text{Cl}_2$  until TLC indicated complete removal of product. Combined organic extracts were concentrated under vacuum to produce 41.0 g of dark brown oil, which was used directly in the next step.


- 53 -

**Step 2**

To a solution of 32.5 g (256 mmol) of crude malondialdehyde 66 in 650 ml of absolute ethanol was added 24.5 g (256 mmol) of guanidine hydrochloride and 17.4 g (256 mmol) of sodium ethoxide. The reaction mixture was refluxed for 4 h, cooled down to room temperature, concentrated and dry loaded on silica gel under vacuum. Flash chromatography (0-10% MeOH/ 20% acetone/ CH<sub>2</sub>Cl<sub>2</sub>) afforded 11.0 g (89.4 mmol; 23% from malonic acid (2 steps)) of pyrimidine 67 as a light yellow solid.


**Step 3**

To a mixture of 166 mg (1.35 mmol) of aminopyrimidine 67, 17 mg (0.14 mmol) of DMAP and 418  $\mu$ L (3.00 mmol) of Et<sub>3</sub>N in 10 mL of THF was added 589 mg (2.7 mmol) of (BOC)<sub>2</sub>O. The mixture was stirred at room temperature for 5 h, concentrated-dry loaded on silica gel and flash chromatographed (1-3% acetone/ CH<sub>2</sub>Cl<sub>2</sub>) to produce 117 mg (0.36 mmol; 27%) of 68 as a clear oil.


**Step 4**

- 54 -

To a solution of 117 mg (0.36 mmol) of aldehyde **68** in 7 mL of  $\text{CH}_2\text{Cl}_2$  was added 67  $\mu\text{L}$  (0.43 mmol) of ethyl isonipecotate and 5  $\mu\text{L}$  of acetic acid. 30 min. later 153 mg (0.72 mmol) of  $\text{NaBH}(\text{OAc})_3$  was introduced. The mixture was stirred overnight at room temperature, diluted with  $\text{CH}_2\text{Cl}_2$ , washed with aqueous  $\text{NaHCO}_3$ , dried and concentrated, and crude residue was flash chromatographed (0-4% sat.  $\text{NH}_3$  in  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) to produce 133 mg (0.29 mmol; 81%) of **69** as a white film.

**Step 5**

To a solution of ester **69** in 5 mL of a 3 : 1 : 1 mixture of THF – water – methanol was added 11 mg (0.44 mmol) of  $\text{LiOH}$ . Reaction mixture was stirred overnight at room temperature, concentrated to dryness and exposed to high vacuum to obtain 134 mg of crude acid **70** as a yellowish solid which was used without purification.

**Example 14****Step 1**


To a  $-78^\circ\text{C}$  solution of 2.36 g (11.4 mmol) of picoline **1A** in 70 mL of THF was added 16.3 mL of 1.4 M  $\text{BuLi}$  solution (22.8 mmol) in hexanes in portions over 10 min.

- 55 -

Reaction mixture was then allowed to warm up and was then stirred for 2 h at room temperature, which resulted in the formation of an orange precipitate. The mixture was cooled back to  $-78^{\circ}\text{C}$ , and ethylene oxide was bubbled through the solution for 1 min. followed by stirring for 5 min. This two-step sequence was repeated eight times.

- 5 The mixture was then allowed to warm to  $-50^{\circ}\text{C}$ , stirred at that temperature for 40 min., quenched with 1.34 mL (23 mmol) of AcOH and allowed to warm to room temperature. Dilution with water was followed by extraction with EtOAc, concentration of the organic phase, and flash chromatography of the crude residue (10-15% acetone/  $\text{CH}_2\text{Cl}_2$  ) to produce 1.50 g (5.95 mmol; 53%) of **71** as a white solid.


10

**Step 2**


- To a  $-60^{\circ}\text{C}$  solution of 628  $\mu\text{L}$  (7.2 mmol) of oxalyl chloride in 20 mL of  $\text{CH}_2\text{Cl}_2$  was added dropwise 1.03 mL (14.5 mmol) of DMSO. After stirring the mixture for 15 min. at  $-55^{\circ}\text{C}$ , a solution of 1.50 g (5.95 mmol) of alcohol **71** in 20 mL of  $\text{CH}_2\text{Cl}_2$  was introduced over the period of 15 min. After the addition was complete, the mixture was stirred for 30 min. at  $-55^{\circ}\text{C}$ , followed by the addition of 4.18 mL (30.0 mmol) of  $\text{Et}_3\text{N}$  and stirring for another 15 min. The reaction mixture was then warmed to room temperature and diluted with water. Extraction with  $\text{CH}_2\text{Cl}_2$  was followed by
- 15
- 20 concentration of the organic phase and flash chromatography (1-15% acetone/  $\text{CH}_2\text{Cl}_2$  ) to produce 1.00 g (4.00 mmol; 67%) of **72** as an off-white solid.



- 56 -


**Step 3**

To a solution of 1.00 g (4.0 mmol) of aldehyde **72** in 25 mL of CH<sub>2</sub>Cl<sub>2</sub> was added 617  $\mu$ L (4.8 mmol) of ethyl isonipecotate followed by one drop of AcOH. Reaction mixture was then stirred for 40 min at room temperature after which 1.70 g (8.0 mmol) of NaBH(OAc)<sub>3</sub> was introduced. Reaction mixture was stirred overnight at room temperature, neutralized with saturated aqueous NaHCO<sub>3</sub>, diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. Concentration and flash chromatography (0 – 4% saturated NH<sub>3</sub> in MeOH/ CH<sub>2</sub>Cl<sub>2</sub>) provided 1.41 g (3.6 mmol; 90%) of **73** as a white solid.

**Step 4**


To a solution of 534 mg (1.47 mmol) of ester **73** in 4 mL of a 3 : 1 : 1 mixture of THF – water – methanol was added 60 mg (2.50 mmol) of LiOH. Reaction mixture was stirred overnight at room temperature, concentrated to dryness and exposed to high vacuum to obtain 540 mg of crude acid **74** as a white solid which was used without purification.

- 57 -

**Example 15**


In a manner similar to that described in Example 6, steps 5, 6, and 7, 70 was converted to 75.

5

**Example 16**

In a manner similar to that described in Example 6, steps 5, 6, and 7, 74 was converted to 76.

10


**Example 17****Step 1**

To a solution of 77 (0.73 g, 3.82 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was added  $(\text{COCl})_2$  (0.41 mL, 4.58 mmol) followed by DMF (0.1 mL) and the reaction was maintained at 40°C for 3 h. The reaction was then concentrated to give a brown solid which was dissolved in  $\text{CH}_2\text{Cl}_2$  (10 mL). N,O-dimethylhydroxylamine hydrochloride (0.56 g, 5.73


15

- 58 -

mmol) and DIPEA (1.33 mL) were added and the reaction was stirred at room temperature overnight. The reaction was quenched by the addition of saturated aqueous  $\text{NaHCO}_3$  and extracted with EtOAc. The combined organic layers were dried and concentrated, and the residue purified by chromatography to give **78** (3.2 g, 84%).


**Step 2**

In a manner similar to that described in Example 5, steps 1 and 4, **78** (0.57 g, 2.41 mmol) was converted to **79** (0.59 g).


**Step 3**

In a manner similar to that described in Example 6, steps 5, 6 and 7, **79** (0.38 g, 1.49 mmol) was converted to **80** (0.24 g).

- 59 -


**Example 18****Step 1**

In a manner similar to that described in Example 6, step 7, **81** (0.36 g, 0.53 mmol; synthesized in the same manner as compound **30**) was converted to **82** (0.34 g, 63%).


**Step 2**

To a solution of **82** (0.115 g, 0.25 mmol) in DMF (4 mL) was added NaH (60% dispersion in mineral oil, 0.03g, 0.76 mmol). After 5 h at room temperature,  $\text{CF}_3\text{CH}_2\text{OSO}_2\text{CF}_3$  (0.069 g, 0.3 mmol) was added and the reaction stirred at room temperature overnight. The reaction was diluted with EtOAc and extracted 3 times with water to remove the DMF. The organic layer was dried and concentrated to give a residue which was purified by chromatography (10% MeOH/ $\text{NH}_3$  in EtOAc) to give **83** (0.08 g, 30%).


- 60 -

**Example 19****Step 1**

To a solution of 17 (0.21 mole, 100ml THF, -10°C) was added 84 (0.14 mole) over 5 min and the reaction mixture became very viscous. Additional THF (100ml) was added and the yellow suspension was warmed from -10°C to 10°C over about 2.5hr. The reaction was quenched by the addition of 100ml saturated NH<sub>4</sub>Cl and 100 ml H<sub>2</sub>O. Extracted once with EtOAc (300ml) and eight times with CH<sub>2</sub>Cl<sub>2</sub> (150 ml). Dried over solid MgSO<sub>4</sub> and filtered. Concentrated and flashed over silica gel chromatography (3 to 10% MeOH (NH<sub>3</sub>)/CH<sub>2</sub>Cl<sub>2</sub>) to obtain 85 (11g, yield: 38%).


**Step 2**

To the mixture of 85 (9.2 g) and MnO<sub>2</sub> (42 g) was added 200ml CH<sub>2</sub>Cl<sub>2</sub>, and the mixture was stirred at room temperature overnight. Additional MnO<sub>2</sub> (20g) was added and the reaction was stirred another 24hrs. The MnO<sub>2</sub> was filtered off and the reaction was concentrated and flashed over silica gel (5% and 10% MeOH (NH<sub>3</sub>)/CH<sub>2</sub>Cl<sub>2</sub>) to give 86 (3.1g, yield: 33%).


**Step 3**

In a manner similar to that described in Example 7, step 2, 86 (3.1g) was converted to 87 (2.0 g, yield: 68%).


- 61 -

**Step 4**

In a manner similar to that described in Example 7, step 3, 4, 5, and 6, 87 was converted to 88.


**Example 20****Step 1**

To the solution of compound 89 in  $\text{CH}_2\text{Cl}_2$  (20ml) at  $0^\circ\text{C}$  was added m-CPBA (0.54g) and the reaction was stirred at  $0^\circ\text{C}$  for 25 min. and then at room temperature stirred for 2 hrs. 40%  $\text{NH}_4\text{OH}$  (12ml) was added and the mixture was stirred for 30min. Separated and extracted the aqueous layer with  $\text{CH}_2\text{Cl}_2$  (10ml). Dried ( $\text{MgSO}_4$ ), filtered and concentrated in vacuo. Flash chromatography (5%  $\text{MeOH}(\text{NH}_3)/\text{CH}_2\text{Cl}_2$ ) gave 90 (0.67g, 80%).

**Step 2**


To the solution of 90 (0.65g) in  $\text{CH}_2\text{Cl}_2$  (6 ml) at  $-10^\circ\text{C}$  was added TFA (6ml) and the reaction was stirred for 1hr from  $-10^\circ\text{C}$  to  $0^\circ\text{C}$ . Concentrated down and azeotroped twice with toluene (20ml), and concentrated to dryness to obtain 91 as a gummy oil which was used as is.

- 62 -

**Step 3**

In a manner similar to that described in Example 7, steps 5 and 6, **91** was converted to **92**.

5


**Example 21****Step 1**

To a solution of **93** (5.17 g, 22.7 mmol) in THF (100 mL) at  $-50^{\circ}\text{C}$  was added *s*-BuLi (38.4 mL of a 1.3M solution in hexane, 49.9 mmol) dropwise. After 1.5h at  $-40^{\circ}\text{C}$ , the reaction was recooled to  $-50^{\circ}\text{C}$  and **95** (4.84 g, 22.7 mmol) in THF (20 mL) was added. After 2.75 h at  $-50^{\circ}\text{C}$ , glacial acetic acid was added followed by saturated aqueous  $\text{NH}_4\text{Cl}$ . The mixture was warmed to room temperature and the layers were separated. The aqueous layer was extracted with EtOAc. The combined organic layers were dried ( $\text{MgSO}_4$ ) filtered and concentrated to give a residue that was purified by flash column chromatography (1% to 3% MeOH/ $\text{NH}_3$  in  $\text{CH}_2\text{Cl}_2$ ) to give **95** (6.35 g, 63%).


10

15


- 63 -

**Step 2**

In a manner similar to that described in Example 12, step 3, **95** (5.34 g, 12.11 mmol) was converted to **96** (4.71 g, 75%).

**Step 3**


In a manner similar to that described in Example 6, step 4, **96** (3.7 g, 8.43 mmol) was converted to **97** (3.08 g, >100%) which was used as is in the next step.

**Step 4**

Compound **97** (0.7 g, 2.25 mmol),  $\text{H}_2\text{NOCH}_3 \cdot \text{HCl}$  (0.94 g, 11.23 mmol) and  $\text{NaOAc}$  (1.47 g, 17.97 mmol) were combined in 1-pentanol (20 mL) and water (2 mL) and heated to reflux for 2 days. The reaction was cooled to room temperature and 0.5 N  $\text{NaOH}$  was added. The  $\text{EtOH}$  was removed in vacuo, additional water (15 mL) was added, and the reaction extracted with 10%  $\text{EtOH}$  in  $\text{CH}_2\text{Cl}_2$  (180 mL total volume). The combined organic extracts were dried and concentrated to give **98** (0.55 g, 92%).




- 64 -

**Step 5**

In a manner similar to that described in Example 6, steps 5, 6, and 7, 98 was converted to 99.

5

**Example 22****Step 1**


prepared according to:  
J. Org. Chem., 1968,  
33(6), 2388

A solution of 2.2 g (9.5 mmol) of 100 in 75 mL of glacial acetic acid was hydrogenated in the presence of 0.5 g of 10% w/w platinum-on-charcoal for 5 h. The reaction mixture was filtered to remove the catalyst and the filtrate was concentrated by evaporation under reduced pressure to produce a solid residue which was basified with 0.5N NaOH and extracted with methylene chloride ( $\text{CH}_2\text{Cl}_2$ ). Methylene chloride extracts were dried over anhydrous  $\text{MgSO}_4$  and concentrated. The residue was purified by flash chromatography eluted with 10 - 30% of 7N  $\text{NH}_3$ -MeOH in  $\text{CH}_2\text{Cl}_2$  to give 0.82 g of 101 (mp 158-163  $^\circ\text{C}$ ). LCMS  $m/z$  240 ( $\text{MH}^+$ ).

10


15

- 65 -

**Step 2**


A mixture of 0.12 g (0.52 mmol) of **101**, 0.2 g (0.52 mmol) of **5A**, 0.67 g (0.5 mmol) of 1-hydroxybenzotriazole hydrate (HOBt), and 0.11 g (0.57 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (DEC) in 7 mL of anhydrous dimethylformamide (DMF) was stirred at ambient temperature for 18 h. The mixture was diluted with water and the resulting precipitate was filtered to produce 0.26 g of **102** as a white solid (mp 110-115 °C). LCMS  $m/z$  557 (MH<sup>+</sup>).

- 66 -

**Step 3**

To a stirred solution of 0.34 g (2.7 mmol) of oxalyl chloride in 3 mL of anhydrous  $\text{CH}_2\text{Cl}_2$  at  $-70^\circ\text{C}$  was added 0.44 g (5.7 mmol) of anhydrous methylsulfoxide in 2 mL of  $\text{CH}_2\text{Cl}_2$ . After being stirred at  $-70^\circ\text{C}$  for 10 minutes, the reaction mixture was added 1.2 g (2.15 mmol) of 102 in 10 mL of  $\text{CH}_2\text{Cl}_2$ . The stirred mixture was kept at  $-70^\circ\text{C}$  for 0.5 h, mixed with 1.8 mL (13 mmol) of triethylamine, and then allowed to warm up to ambient temperature by itself. The mixture was diluted with water and extracted with  $\text{CH}_2\text{Cl}_2$ . Organic extracts were washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated to produce 1.18 g of 103 as a glass. LCMS  $m/z$  555 ( $\text{MH}^+$ ).

- 67 -

**Step 4**


A solution of 0.8 g (1.44 mmol) of **103** and 0.6 g (7.2 mmol) of methoxylamine hydrochloride in 40 mL of ethanol and 40 mL of pyridine was heated under reflux for 18 h. The mixture was concentrated and the residue was taken up in ethyl acetate/ether and washed with water. The organic solution was dried over anhydrous MgSO<sub>4</sub> and concentrated to 0.65 g of viscous residue which was dissolved in 8 mL of trifluoroacetic acid and 8 mL of CH<sub>2</sub>Cl<sub>2</sub> and stirred at ambient temperature for 18 h. The solution was concentrated and the residue was basified with 1N NaHCO<sub>3</sub> and extracted with ethyl acetate. Organic extracts were washed with brine, dried over anhydrous MgSO<sub>4</sub> and concentrated to a gummy residue. Purification of this residue by flash chromatography with 5 - 8% of 7N NH<sub>3</sub>-MeOH in CH<sub>2</sub>Cl<sub>2</sub> produced 0.151 g of **104** as a gum, LCMS m/z 484 (MH<sup>+</sup>) and 0.146 g of **105** as a glass, LCMS m/z 556 (MH<sup>+</sup>).

- 68 -

Mixing a solution of 0.056 g of the free base of **104** in ethyl acetate with a solution of 0.04 g of maleic acid in ethyl acetate produced a precipitate which was isolated by filtration to give 0.06 g of a dimaleate salt of **104** (mp 155-160 °C).

5 **Example 23**

**Step 1**




J. Med. Chem., 1976,  
19, 360

2.4 g (10. mmol) of **106** were reduced in the similar manner as that described in Example 22, step 1 to give 1.5 g of **107** as a semi-solid. LCMS m/z 240 (MH<sup>+</sup>).

10


**Step 2**



1.5 g (6.31 mmol) of **107** were coupled with **3** in the similar manner as that described in Example 22, step 2 to give 3-g of **108** as a solid ( mp 104-106 °C).


15 LCMS m/z 557 (MH<sup>+</sup>).

- 69 -


**Step 3**

1.17 g (2.1 mmol) of 108 were oxidized in the similar manner as that described in Example 22, step 3 to give 0.7g of 109 as a glass. LCMS  $m/z$  557 ( $MH^+$ ).

- 70 -

**Step 4**


0.32 g (0.58 mmol) of 109 were reacted with 0.6 g (7.2 mmol) of methoxylamine hydrochloride in the same manner as that described in Example 22, step 4 to provide 0.065 g of 110 as a gum, LCMS  $m/z$  484 (MH<sup>+</sup>) and 0.12 g of 111 as a glass, LCMS  $m/z$  556 (MH<sup>+</sup>).

**Example 24****Step 1**


J. Med. Chem., 1994,  
37 (16), 2537

- 71 -

A mixture of 18 g (74 mmol) of **112**, 7.2 g (74 mmol) of N,O-dimethylhydroxylamine hydrochloride, 19.4 g (15 mmol) of N,N-diisopropylethylamine, 1.1 g (8 mmol) of HOBt and 14.2 g (74 mmol) of DEC in 80 mL of anhydrous DMF was stirred at ambient temperature for 18 h. The mixture was diluted with water and extracted with ethyl acetate. Organic extracts were washed with 1% NaHCO<sub>3</sub> and brine, dried over anhydrous MgSO<sub>4</sub> and concentrated to give 15.5 g of **113** as an oil. LCMS m/z 287 (MH<sup>+</sup>).

**Step 2**


To a stirred solution of 2.9 g (18 mmol) of 2-bromopyridine in 30 mL of anhydrous THF at -78°C was added 7.5 mL of 2.5M solution of n-BuLi in hexane dropwise for 0.5 h. After being stirred at -78°C for 1 h, the reaction mixture was added a solution of 5.1 g (17.8 mmol) of **113** in 15 mL of THF. The mixture was allowed to stir at ambient temperature for 48 h, mixed with saturated aqueous NH<sub>4</sub>Cl and extracted with ether. Organic extracts were washed with brine, dried over anhydrous MgSO<sub>4</sub> and concentrated to produce 5.7 g of **114** as an oil. LCMS m/z 305 (MH<sup>+</sup>).

**Step 3**


A solution of 3.15 g (10.4 mmol) of **114** and 3.47 g (41.6 mmol) of methoxylamine hydrochloride in 30 mL of ethanol and 30 mL of pyridine was heated under reflux for 18 h. The mixture was concentrated and the residue was taken up in ether and washed with water. The organic solution was dried over anhydrous MgSO<sub>4</sub> and concentrated to give 2.5 g of **115** as an oil. LCMS m/z 334 (MH<sup>+</sup>).



- 72 -


**Step 4**

A solution of 2.4 g (7.2 mmol) of 22 in 20 mL of  $\text{CH}_2\text{Cl}_2$  and 20 mL of trifluoroacetic acid was stirred at ambient temperature for 1 h. The solution was concentrated. The residue was basified with saturated aqueous  $\text{NaHCO}_3$  and extracted with  $\text{CH}_2\text{Cl}_2$ . Organic extracts were washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated to give 1.41 g of 23 as a glass. LCMS  $m/z$  234 ( $\text{MH}^+$ ).


**Step 5**

A mixture of 0.466 g (2 mmol) of 116, 0.517 g (2.2 mmol) of 5A, 0.276 g (2 mmol) of HOBt and 0.46 g (2.4 mmol) of DEC in 20 mL of anhydrous DMF was stirred at ambient temperature for 18 h. The mixture was concentrated by evaporation under reduced pressure at bath temperature of 25-45°C and the residue was chromatographed with 4% (7N  $\text{NH}_3/\text{CH}_3\text{OH}$ ) in  $\text{CH}_2\text{Cl}_2$  to produce 0.48 g of syrup which was dissolved in 15 mL of EtAc-EtOH (3:1 v) and mixed with a solution of 0.26 g of maleic acid in 10 mL of EtAc-EtOH (1:1). The resulting precipitate was filtered to produce 0.35 g of the maleate salt of 117 (mp 160-163 °C). LCMS  $m/z$  451 ( $\text{MH}^+$ ).

- 73 -


**Example 25****Step 1**

To a stirred solution of 4.16 g (20 mmol) of 1A in 80 mL of anhydrous THF at  
 5 -78°C was added dropwise 17 mL of 2.5M solution of n-BuLi in hexane for 25  
 minutes. After being stirred from -78°C to room temperature for 1 h, the reaction  
 mixture was added a solution of 6 g (22 mmol) of 26 in 100 mL of anhydrous THF and  
 kept at room temperature for 18 h. The mixture was mixed with saturated aqueous  
 NH<sub>4</sub>Cl and extracted with EtAc. Organic extracts were washed with brine, dried over  
 10 anhydrous MgSO<sub>4</sub> and concentrated to produce 6.1 g of 118 (mp 146-149 °C). LCMS  
 m/z 420 (MH<sup>+</sup>).


**Step 2**

15 A solution of 3.71 g (8.8 mmol) of 118 and 3.7 g (44 mmol) of methoxylamine  
 hydrochloride in 40 mL of pyridine and 40 mL of ethanol was heated under reflux for 2  
 days. The mixture was concentrated and the residue was taken up in CH<sub>2</sub>Cl<sub>2</sub> and  
 washed with saturated aqueous NaCl. Organic solution was dried over anhydrous  
 MgSO<sub>4</sub> and concentrated to give 2.6 g of 119 as a glass. LCMS m/z 421 (MH<sup>+</sup>).


- 74 -

**Step 3**

A solution of 0.9 g (2.14 mmol) of **119** in 10 mL of  $\text{CH}_2\text{Cl}_2$  and 10 mL of trifluoroacetic acid was stirred at ambient temperature for 2 h. The solution was concentrated. The residue was taken up in  $\text{CH}_2\text{Cl}_2$ , washed with saturated  $\text{NaHCO}_3$  and brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated to a solid residue which was triturated with  $\text{CH}_3\text{CN}$  and filtered to produce 0.29 g of **120** (mp 200-205 °C). LCMS  $m/z$  321 ( $\text{MH}^+$ ).


**Step 4**

0.1 g (0.31 mmol) of **120** and 0.83 g (0.35) of **5A** were coupled in the same manner as that described in Example 24, step 5 to produce 0.12 g of the maleate salt of **121** (mp 170-173 °C). LCMS  $m/z$  538 ( $\text{MH}^+$ ).



**Example 26****Step 1**

In a similar manner to that described in Example 6, step 7, **122** (0.26 g, 0.41 mmol) was converted to **123** (0.08 g, 40%).

- 75 -


**Example 27****Step 1**

To a suspension of LAH (0.83 g, 22 mmol) in ether (20 mL) at 0° C was added  
5 **124** (3.2 g, 17.5 mmol) in THF (15 mL) dropwise. The reaction was stirred at 0° C for  
1.5 h, and quenched by the addition of water (0.8 mL), 20% aqueous NaOH (0.8 mL),  
and water (2.4 mL). The mixture was stirred for 15 min and filtered and the filter cake  
washed with CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was concentrated to give an oil which was  
dissolved in ether (30 mL) and washed with brine and dried (MgSO<sub>4</sub>). Filtration and  
10 concentration in vacuo gave **125** (2.5 g) which was used without further purification.

**Step 2****Step 3**


In a similar manner to that described in Example 22, step 3 and Example 1,  
steps 4, 5, and 6, **125** was converted to **126**.

- 76 -

**Step 4**

In a similar manner to that described in Example 6, step 5, 126 was converted to 127.

5











**Step 5**











In a similar manner to that described in Example 6, step 7, 127 was converted to 128.













10

The compounds in Table 1 (first column) are prepared from the compounds in the last column of Table 1 by following essentially the same procedures as in the examples described above. In Table 1 "Cmpd. No." stands for "Compound Number".













TABLE 1













Cmpd. No.	STRUCTURE	Mass Spec. $[M^+H]^+$	Starting Material
200		470.1	 +HCl
201		456.1	 +HCl
202		456.1	 +HCl
203		531.1	 +HCl
204		499.1	 +HCl













205		497.1	
206		517.1	
207		549.1	
208		599.1	
209		568.1	













210		565.1	 ·HCl
211		483	 ·HCl
212		484.1	 ·HCl
213		583.1	 ·HCl
214		552.1	 ·HCl
215		471	 ·HCl



























216		512	 ·HCl
217		512	
218		504	
219		454	
220		470	
221		456	 ·HCl













222		456	 ·HCl
223		495	
224		470	 ·HCl
225		470	 ·HCl
226		504	
227		484	 ·HCl















228		472	
229		486	
230		572	
231		505	
232		452	
233		518	

234		450	
235		442	
236		423	
237		423	
238		436	
239		451	

















240		423	
241		423	
244		435	
245		519	
246		451	
247		421	











248		438	
249		452	
250		487	
251		543	
252		501	
253		457	

254		471	
255		465	
256		465	
257		422	
258		406	
259		455	

260		484	
261		443	
262		440	
263		441	
264		427	
265		427	
266		518	





267		490	 ·HCl
268		455	 ·HCl
269		439	 ·HCl
270		407	 ·HCl
271		421	 ·HCl
272		407	 ·HCl
273		455	 ·HCl
275		425	 ·HCl

278		425	 +HCl
279		439	 +HCl
280		470	 +HCl
281		469	 +HCl
282		504	

- 90 -


The isomers 246A and 253A, below, can be separated from 246 and 253, respectively, above, by techniques well known to those skilled in the art.



5

### Example 28

#### Step 1



285

To a solution of 1.00 g (8.13 mmol) of pyrimidine aldehyde **67** (Step 2 of Example 13) in 40 ml of  $\text{CH}_2\text{Cl}_2$  was added 1.36 mL (10.58 mmol) of ethyl isonipecotate and 2 drops of acetic acid. The mixture was stirred for 40 min. at room temperature, after which 2.58 g (12.17 mmol) of  $\text{NaBH}(\text{OAc})_3$  was added. The reaction mixture was then stirred for 20 h at room temperature, diluted with aqueous NaOH (pH adjusted to 11) and extracted with  $\text{CH}_2\text{Cl}_2$ . Organic phase was dried and concentrated, and the residue was flash chromatographed (4-8% ca. 3 N  $\text{NH}_3$  in


10

15

- 91 -

MeOH/ CH<sub>2</sub>Cl<sub>2</sub>) to produce 1.55 g (5.87 mmol; 72%) of amine 285 as a yellowish solid.


### Step 2



286

To a solution of 3.83 g (14.51 mmol) of ester 285 in 60 ml of 3 : 1 : 1 mixture of THF – MeOH – H<sub>2</sub>O was added 1.22 g (29.02 mmol) of LiOH monohydrate. The reaction mixture was stirred at room temperature overnight, concentrated, and the residue was dried under high vacuum to produce 3.84 g of crude acid 286 lithium salt as a yellow solid. Material could be used directly or could be purified by passing through a silica gel plug eluting with ca. 3 N NH<sub>3</sub> in MeOH.

### Step 3



287

To a mixture of 3.32 g (14.05 mmol) of acid 286 and 4.07 g (14.05 mmol) of 4-[(E)-(methoxyimino)-2-pyridinylmethyl]piperidine dihydrochloride (see Compound 447 below) in 40 mL of DMF was added 8.94 mL (70.25 mmol) of 4-ethylmorpholine and 14.0 mL (23.52 mmol) of 50 wt. % solution of 1-propanephosphonic acid cyclic anhydride in ethyl acetate. The reaction mixture was stirred for 4.5 h at 50°C followed by 14 h at room temperature. Concentration of the mixture was followed by exposure to high vacuum for 24 h to remove remaining DMF. The residue was partitioned between aqueous NaOH and CH<sub>2</sub>Cl<sub>2</sub>, organic phase was separated, dried and


- 92 -

concentrated, and the residue was flash chromatographed (5-15% ca. 3 N NH<sub>3</sub> in MeOH/ CH<sub>2</sub>Cl<sub>2</sub>) to produce 4.60 g (10.51 mmol; 75 %) of amide 287 as a light tan foam. MS 438 (M+1).

5 **Example 29**


**Step 1**

Reference: *J. Heterocyclic Chem.*, **1966**, 3, 252.



3,4 Pyridine-dicarboximide 288 (10.0 g; 67.5 mmoles) was dissolved in 162 g.  
10 of 10% aqueous NaOH and the solution was cooled to an internal temperature of 7 °C  
in an ice-salt bath. Bromine (3.6 ml; 70 mmoles) was added dropwise. After the  
addition, the solution was heated for 45 minutes at a bath temperature of 80-85 °C.  
The yellow solution was then cooled to an internal temperature of 37 °C, then 17 ml of  
glacial acetic acid were added dropwise to a pH of 5.5. The resulting mixture was  
15 saved overnight in a refrigerator. The solid formed was filtered and washed with 5 ml  
of water and 5 ml of methanol. The reaction yielded 6.35 g. of product 289 melting at  
280-285 °C (decomp.).


**Step 2**



20 Solid Compound 289 (9.5 gr.; 69 mmoles) was carefully added in three aliquots  
to a slurry of lithium aluminum hydride (9.5 gr.; 250 mmoles) in 200 ml of dry  
tetrahydrofuran. The resulting hot mixture was stirred at room temperature for two  
days. After cooling in an ice bath, the reaction was quenched with very careful  
25 sequential dropwise addition of 10 ml of water, followed by 10 ml of 15% aqueous


NaOH, then by 30 ml of water. The resulting solid was filtered through a pad of Celite and washed several times with THF. The oil obtained after evaporation of the solvent, solidified on standing. The reaction mixture was purified by flash chromatography on silica gel using 5%MeOH(NH<sub>3</sub>)/EtOAc as eluent yielding 6.21 (72%) of Compound 290 . LC-MS: m/z = 125 (M+1).

### Step 3



Manganese dioxide (29 gr.; 334 mmoles) was added, in one portion, at room temperature, to a suspension of 3-amino-4-hydroxymethyl pyridine **290** (5.0 gr.; 40.3 mmoles) in 500 ml of chloroform with good stirring. After two days, the solid is filtered through a pad of Celite and washed with chloroform. Removal of the solvent using reduced pressure yielded 4.2 grams (85%) of Compound **291** as a yellow solid.


### Step 4




A dry dichloromethane (400 ml) solution of ethyl isonipecotate (12.5 gr.; 79.5 mmoles) and 3-amino pyridine 4-carboxyaldehyde **291** (3.33 gr.; 27.3 mmoles) was stirred at room temperature for one hour, then 60 grams of activated 3Å molecular sieves were added. The mixture was stirred for an additional 90 minutes, then 20 grams (96.4 mmoles) of sodium triacetoxo borohydride was added at room temperature in one portion. After stirring for three days, the solid was filtered through a pad of Celite and washed with dichloromethane. The solution was stirred for 15

- 94 -

minutes with 100 ml of saturated aqueous sodium bicarbonate then separated from the aqueous layer. The organic layer was washed two more times with saturated aqueous sodium bicarbonate, then with brine and dried with anhydrous sodium sulfate. After evaporation of the solvent, the resulting oil was purified by flash chromatography on silica gel using EtOAc:Hexanes:MeOH(NH<sub>3</sub>) as eluent. The procedure yielded 6.8 gr.(94%) of Compound 292 . FAB-MS: m/z = 264 (M+1).

**Step 5**


10 Ethyl 1-[(3-amino-4-pyridinyl)methyl]-4-piperidinecarboxylate 292 (4.75 gr.; 18.04 mmoles) was stirred for 24 hours at room temperature with 1.51 gr. (36 mmoles) of lithium hydroxide monohydrate in 75 ml of methanol. Removal of the solvent using reduced pressure yielded Compound 293 as a white solid.

**Step 6**

15 4-(2-pyridinylcarbonyl)piperidine 28 (Step 4 in Example 6) (0.3 gr.; 1.58 mmoles), lithium 1-[(3-amino-4-pyridinyl)methyl]-4-piperidinecarboxylate 293 (0.34 gr.; 1.4 mmoles), DEC (0.38 gr.; 2.0 mmoles), and HOBt (0.27 gr.; 2.0 mmoles) were  
20 stirred at room temperature in 10 ml of dry DMF for two days. The reaction was

- 95 -

quenched with 50 ml. of 0.5 N aqueous NaOH, then the solution was extracted with dichloromethane. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. The product 295 was isolated by flash chromatography on silica gel using EtOAc:Hexanes:MeOH(NH<sub>3</sub>) (50:45:5) as eluent. Yields: 0.27 gr. (47%). FAB-MS: m/z = 408 (M+1).

**Step 7**

1-[[[1-[(3-amino-4-pyridinyl)methyl]-4-piperidinyl]carbonyl]-4-(2-pyridinylcarbonyl)piperidine 295 (0.196 gr.; 0.48) and methoxyamine hydrochloride (0.401 gr. 4.8; mmols) were heated, under N<sub>2</sub>, at a bath temperature of 70°C for 24 hours in 6.0 ml of dry pyridine. After removing the pyridine using reduced pressure, the residue was treated with saturated aqueous sodium bicarbonate. The resulting mixture was extracted several times with dichloromethane. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. The reaction mixture was purified by silica gel preparative thin layer chromatography. The plates were eluted with EtOAc:Hexanes:MeOH(NH<sub>3</sub>) (60:35:5) and the product 296 was extracted with 10% MeOH(NH<sub>3</sub>)/EtOAc. Yields: 0.15 gr. (71%). FAB-MS: m/z = 437 (M+1).


**Example 30 Step 1**

A mixture of 297 (1 g, 10 mmol) in 1:1 water-dioxane (50 mL) was treated with Et<sub>3</sub>N (4 mL, 13 mmol) and BOC<sub>2</sub>O (2.8 g, 13 mmol) at 4°C and allowed to warm to 20°C for one day. The solvent was then removed in vacuo. The residue was taken up in 1:1 water-ethyl acetate and the organic layer was discarded. The aqueous




- 96 -

layer was acidified with 1 N aqueous HCl and extracted three times with ethyl acetate. The combined organic phases were washed with water and brine, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated to give **298** as a white solid (1.8 g, 90%).

5 **Step 2**


A mixture of **298** (1.8 g, 9 mmol), N,O-dimethylhydroxylamine hydrochloride (2.6 g, 27 mmol), EDCI (5 g, 27 mmol), HOBT (0.1 g, 1 mmol), and DIPEA (12.5 mL, 72 mmol) in DMF (30 mL) was stirred at 20°C overnight. The reaction was then  
10 concentrated to half volume in vacuo, poured onto water, and extracted three times with ethyl acetate. The combined organic phases were washed with saturated aqueous  $\text{NH}_4\text{Cl}$ , saturated aqueous  $\text{NaHCO}_3$ , water and brine, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated to give **299** as a clear oil (2.1 g, 98%).

15 **Step 3**

To a solution of 2-bromopyridine (1.2 mL, 12 mmol) in THF (60 mL) at -78°C was added n-BuLi (8 mL of a 1.6 M solution in hexanes, 12 mmol) dropwise over 15 min. After stirring for an additional 30 min at -78 °C, a solution of **299** (1 g, 4 mmol) in  
20 THF (20 mL) was slowly added. The reaction was then heated to 60 °C for 1 h. After cooling to 20 °C, the reaction was diluted with ether, quenched with saturated aqueous  $\text{Na}_2\text{SO}_4$ , and dried with solid  $\text{Na}_2\text{SO}_4$ . The mixture was filtered through a plug of solid  $\text{Na}_2\text{SO}_4$  and concentrated in vacuo. Flash column chromatography (0–20% ethyl acetate–hexanes) yielded **300** as a yellow oil (0.12 g, 11%).

25






- 97 -








**Step 4**








Following procedures similar to those of Steps 4 to 7 of Example 6, compound 301 was obtained. MS 409 (M+1).








5








Following procedures similar to those described in the examples above, the compounds in Table 2 were prepared.

TABLE 2		
Compound	STRUCTURE	MS (M+1)
302		430
303		421
304		505
305		505
306		471

307		426
308		408
309		442
310		437
311		437
312		458
313		402






314		487
315		438
316		467
317		424
318		451
319		430
320		523








321		453
322		453
323		410
324		413
325		439
326		466
327		453








328		453
329		424
330		453
331		438
332		488
333		437
334		437








335	 <chem>CC(C)OCCN=C(c1ccncc1)C2CCN(C2)C(=O)C3CCN(C3)CCc4cc(N)ncn4</chem>	479
336	 <chem>COC(=O)C1CCN(C1)C(=O)C2CCN(C2)CCc3cc(N)ncn3C(CO)C(CO)Cc4cc(N)ncn4</chem>	452
337	 <chem>COC(=O)C1CCN(C1)C(=O)C2CCN(C2)CCc3cc(N)ncn3C(COC)C(COC)Cc4cc(N)ncn4</chem>	466
338	 <chem>COC(=O)C1CCN(C1)C(=O)C2CCN(C2)CCc3cc(N)ncn3C(COC)C(COC)Cc4cc(N)ncn4</chem>	438
339	 <chem>CC(C)OCCN=C(c1ccncc1)C2CCN(C2)C(=O)C3CCN(C3)CCc4cc(N)ncn4</chem>	465
340	 <chem>CCOC(=O)C1CCN(C1)C(=O)C2CCN(C2)CCc3cc(N)ncn3C(COC)C(COC)Cc4cc(N)ncn4</chem>	465













341		513
342		452
343		550
344		499
345		451




346		451
347		451
348		451
349		452
350		455
351		455
352		422

353		422
354		492
355		438
356		437
357		424
358		510
359		539

360		453
361		409
362		438
363		426
364		422
365		483
366		483

367		497
368		465
369		479
370		479
371		493

372		564
373		517
374		568
375		426
376		455















377	 <chem>CO=Nc1cc(F)ccn1C2CCN(C2)C(=O)C3CCN(C3)Cc4cc(N)ccn4</chem>	456
378	 <chem>CO=Nc1cc(C)ccn1C2CCN(C2)C(=O)C3CCN(C3)Cc4cc(N)ccn4</chem>	452
379	 <chem>O=Cc1cc(F)ccn1C2CCN(C2)C(=O)C3CCN(C3)Cc4cc(N)ccn4</chem>	427















- 111 -















If one were to follow procedures similar to those described in the examples above, the compounds in the "Structure" column of Table 3 would be obtained using the starting material listed in Table 3. Each compound in Table 3 is a mixture of oxime isomers, as represented by the  $\sim$  bond between the oxime nitrogen and the OH or OCH<sub>3</sub> moiety. In Table 3 "CMPD" stands for "Compound".































TABLE 3















CMPD	Structure	Starting Material
380		 •HCl
381		 •HCl
382		 •HCl
383		 •HCl
384		
385		 •HCl
386		















CMPD	Structure	Starting Material
387		
388		
389		
390		
391		
392		
393		















CMPD	Structure	Starting Material
394		
395		
396		
397		
398		
399		
400		

CMPD	Structure	Starting Material
401		
402		
403		
404		
405		
406		
407		









CMPD	Structure	Starting Material
408		
409		
410		
411		
412		
413		
414		

CPD	Structure	Starting Material
415		
416		
417		
418		
419		
420		
421		


CMPD	Structure	Starting Material
422		
423		
424		
425		
426		
427		
428		

CMPD	Structure	Starting Material
429		
430		
431		
432		
433		
434		
435		




CMPD	Structure	Starting Material
436		
437		
438		
439		


- 121 -

**Example 31****Step 1**

5 To a solution of LDA (233 mL, 2.0 M in THF/heptane/ethylbenzene, 0.466 mol) in THF (300mL) at 0 °C was added, dropwise over 1.0 h, a solution of compound **440** (100g, 0.389 mol) in THF (~ 400 mL). The red-orange solution was stirred at 0 °C for 30 min, and then transferred by cannula to a pre-cooled (0 °C) solution of N-fluorobenzenesulfonimide (153 g, 0.485 mol) in dry THF (~ 600 mL). The reaction  
10 mixture was stirred at 0 °C for 30 min, and then at rt for 18 h. The total solvent volume was reduced to approximately one third, and EtOAc (~ 1L) was added. The solution was washed successively with water, 0.1 N aq. HCl, saturated aq. NaHCO<sub>3</sub>, and brine. The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated under reduced pressure to yield a crude liquid. Separation by flash chromatography (6:1  
15 hexanes-EtOAc) gave compound **441** (93.5 g, 87%).

**Step 2**


20 In a manner similar to that described in Example 6, Step 4, compound **441** was converted to compound **442**.

**Step 3**


25

- 122 -

In a manner similar to that described in Example 1, Step 4, compound 442 was converted to compound 443.


**Step 4**

In a manner similar to that described in Example 1, Step 5, compound 443 was converted to compound 444.

**Step 5**

In a manner similar to that described in Example 6, Step 5, compound 5 was converted to compound 445.


15

**Step 6**


In a manner similar to that described in Example 6, Step 6, compound 445 was converted to compound 446.

- 123 -


In the above examples, the compound 4-[(E)-(methoxyimino)-2-pyridinylmethyl]piperidine dihydrochloride:



- 5 can be used to prepare the compounds of this invention, for example, see Examples 6 and 28. Preferably, Compound 447 is prepared from a compound of formula:




and from a compound of Formula 449:



$R^{50}$  is an alkyl or aryl group,  $f$  is 0 to 4,  $R^{51}$  is an alkyl group, and  $Q$  is a halo group, wherein said alkyl, aryl, and halo groups are as defined above.


Compound 447 can be prepared from 448 and 449 by:

- 15 (a) converting the compound of formula 449 into its Grignard form (449A):




- 20 (b) reacting the compound of formula 448 with the compound of formula 449A to obtain a compound of formula 450:

- 124 -




(c) reacting the compound of formula 450 with a suitable alkyl chloroformate of formula 451

5




to yield a compound of formula 452:




10

(d) forming the salt (formula 453):



(e) reacting the compound of formula 453 with an alkoxyamine ( $\text{NH}_2\text{OR}^{51}$ ) or its hydrochloride to form an oxime of formula 454:

15




(f) isomerizing the compound of formula 454 by treatment with a strong acid and simultaneously converting to the desired acid salt of Formula 454 with an enriched E isomer, wherein the E isomer predominates over the Z-isomer by at

20

- 125 -


least a 90:10 ratio. When  $f=0$ ,  $R^{51}$  is methyl, and the acid used for isomerization is HCl in the compound of formula 454, the final product is the compound of formula 447.

This preparation can be represented as follows:



- 126 -

Following the above process the Compound 447 can be prepared as follows:



The conversion of compound 461 to 447 predominantly yields the E-isomer of compound 447 in high stereochemical purity and high yields. Isomerization of a mixture of phenyl compounds by acid catalysis is discussed by T. Zsuzsanna *et al*, *Hung.Magy.Km.Foly.*, 74(3) (1968), 116-119.

The above process starts with Compound 449. In step 1, a 4-halo-1-alkylpiperidine (or a 4-halo-1-arylpiperidine) is converted to its Grignard analog (449A) by reacting with magnesium. The reaction is performed generally at temperatures of about  $-10^{\circ}\text{C}$  to reflux. Generally a hydrocarbon solvent such as, for example, toluene, xylene, chlorobenzene, dichlorobenzene and the like, or mixture of hydrocarbons listed above with an ether, such as, for example, a  $\text{C}_5\text{-C}_{12}$  alkyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, diglyme, 1,4-dioxane, tetrahydrofuran and the like are suitable for this reaction. The solution is cooled to around  $-10^{\circ}\text{C}$  to about  $10^{\circ}\text{C}$  and then reacted with a suitable 2-cyanopyridine (448), for about 10-120 minutes. Examples of suitable 2-cyanopyridines are 2-cyanopyridine, 4-methyl-2-

- 127 -

cyanopyridine, 4-ethyl-2-cyanopyridine, 4-phenyl-2-cyanopyridine, and the like. Preferred are 2-cyanopyridine and 4-methyl-2-cyanopyridine. The Grignard compound is used generally in about 1-4 molar equivalents with respect to the compound of formula 448, preferably in about 1-3 molar equivalents and typically in about 1.5-2.5 molar equivalents. The product of formula 450 may be isolated by procedures well known in the art, such as, for example, treatment with an acid (e.g. HCl); preferably in a suitable solvent (e.g., tetrahydrofuran or ethyl acetate).

The product of Formula 450 may then be reacted with an alkyl chloroformate in the next step. Suitable alkyl chloroformates are, for example, methyl chloroformate, ethyl chloroformate, propyl chloroformate, and the like, with the preferred being methyl chloroformate or ethyl chloroformate. Generally a hydrocarbon solvent such as, for example, toluene, xylene, chlorobenzene, dichlorobenzene and the like, or mixture of a hydrocarbons listed above with an ether such as, for example, a C<sub>5</sub>-C<sub>12</sub> alkyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, diglyme, 1,4-dioxane, tetrahydrofuran and the like is suitable for this reaction. The reaction is generally performed at about 25-100°C, preferably about 40-90°C and typically about 50-80°C, for about 1-5 hours. After the reaction, generally the generated acid is washed off and the product of formula 452 may be isolated by organic solvent extraction.

The compound of Formula 452 may then be converted into its acid salt by treatment with an acid such as, for example, sulfuric acid, hydrochloric acid, trifluoroacetic acid and the like, generally in a solvent at temperatures between ambient and reflux of the solvent. Suitable solvents include hydrocarbons such as, for example, toluene, xylene, chlorobenzene, dichlorobenzene and the like. There being two nitrogen atoms in the compound of Formula 452, the salt generally has 2 moles of acid to a mole of compound 452.

The compound of formula 453 may then be converted to an alkyloxime of formula 454 by reacting it with an alkoxyamine (or its hydrochloride), usually in aqueous solution form. Suitable alkoxyamines are, for example, methoxyamine, ethoxyamine and the like. Methoxyamine is preferred. The alkoxyamine (or its hydrochloride) is employed generally in about 1 to about 4 molar equivalents, preferably in about 1 to about 3 molar equivalents, and typically in about 1 to about 2 molar equivalents. Generally, the reaction is catalyzed by a weak acid such as, for example, acetic acid, formic acid and the like, or mixtures thereof. A cosolvent such



- 128 -

as, for example, methanol, ethanol, isopropanol, n-butanol and the like, or mixtures thereof may be added. The product of formula 454, after work-up, is a mixture of the Z- and the E-isomers, whose ratio may be analyzed for its stereochemical make-up, using techniques well known in the art such as, for example, HPLC.

5 Treating the compound of formula 454 with a strong acid under the reaction conditions described below isomerizes the mixture of the Z and the E-isomers into predominantly the E-isomer. Generally, the compound of formula 454 may be dissolved in a solvent such as, for example, ethanol, methanol, isopropanol, n-butanol and the like, ether such as methyl tert-butyl ether, tetrahydrofuran and the like,  
10 hydrocarbon such as, for example, heptane, hexane, toluene and the like, nitrile such as, for example, acetonitrile, benzonitrile and the like, or mixtures of such solvents. The dissolved compound is then treated with a strong acid such as, for example, HCl, HBr, H<sub>2</sub>SO<sub>4</sub> and the like, at temperatures in the range of 20 to 100°C for about 1-20 hours. The acid is employed generally in about 1 to about 8 molar equivalents,  
15 preferably in about 1 to about 6 molar equivalents, and typically in about 2 to about 4 molar equivalents. Work-up typically forms predominantly the acid salt of the E-isomer of the compound of formula 454, which is in fact the compound of formula 447 when R<sup>51</sup> = methyl, n=0 and the acid salt is HCl in 454.

The products of the various steps in the process described above may be  
20 isolated and purified by conventional techniques such as, for example, filtration, recrystallization, solvent extraction, distillation, precipitation, sublimation and the like, as is well known to those skilled in the art. The products may be analyzed and/or checked for purity by conventional methods such as, for example, thin layer chromatography, NMR, HPLC, melting point, mass spectral analysis, elemental  
25 analysis and the like, well known to those skilled in the art.

### H<sub>3</sub>-Receptor Binding Assay

The source of the H<sub>3</sub> receptors in this experiment was guinea pig brain. The animals weighed 400-600 g. The brain tissue was homogenized with a solution of 50  
30 mM Tris, pH 7.5. The final concentration of tissue in the homogenization buffer was 10% w/v. The homogenates were centrifuged at 1,000 x g for 10 min. in order to remove clumps of tissue and debris. The resulting supernatants were then centrifuged at 50,000 x g for 20 min. in order to sediment the membranes, which were

- 129 -

next washed three times in homogenization buffer (50,000 x g for 20 min. each). The membranes were frozen and stored at -70°C until needed.

All compounds to be tested were dissolved in DMSO and then diluted into the binding buffer (50 mM Tris, pH 7.5) such that the final concentration was 2µg/ml with 0.1% DMSO. Membranes were then added (400 µg of protein) to the reaction tubes.

The reaction was started by the addition of 3 nM [<sup>3</sup>H]R-α-methyl histamine (8.8

Ci/mmol) or 3 nM [<sup>3</sup>H]N<sup>α</sup>-methyl histamine (80 Ci/mmol) and continued under incubation at 30°C for 30 min. Bound ligand was separated from unbound ligand by filtration, and the amount of radioactive ligand bound to the membranes was

quantitated by liquid scintillation spectrometry. All incubations were performed in duplicate and the standard error was always less than 10%. Compounds that inhibited more than 70% of the specific binding of radioactive ligand to the receptor were serially diluted to determine a K<sub>i</sub> (nM).

Compounds 23, 30, 31, 32, 33, 44, 45, 49, 50, 53, 54, 55, 56, 57A, 59, 75, 76, 83, 88, 92, 99, 104, 110, 117, 128, 200, 201, 203-215, 217-241, 244-246, 246A, 247-253, 253A, 254-273, 275, 278, 280-282, 287, 296, 301-310, and 312-379 had a K<sub>i</sub> within the range of about 0.25 to about 370nM.

Preferred Compounds 23, 30, 31, 32, 33, 50, 53, 54, 55, 56, 57A, 59, 92, 212, 215, 218, 219, 220, 224, 225, 226, 227, 229, 233, 235, 237, 238, 246, 246A, 247, 248, 251, 253, 253A, 268-273, 275, 278-281, 287, 296, 301, 304-307, 309, 312, 314-318, 320-356, and 358-376 had a K<sub>i</sub> within the range of about 0.25 to about 33nM.

Most preferred Compounds 30, 31, 32, 33, 54, 55, 56, 56A, 225, 237, 246A, 253A, 273, 280, 287, 296, 301, 304-307, 309, 312, 314-318, 320-348, 350-356, 359-372, and 374-376 had a K<sub>i</sub> within the range of about 0.25 to about 16nM.

More preferred compound 32 had a K<sub>i</sub> of 0.83nM.

More preferred compounds 54, 55, 253A, 287, 320 had a K<sub>i</sub> within the range of about 1.05 to about 9.75nM.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid.

Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art,

- 130 -

e.g. magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), Remington's  
5 Pharmaceutical Sciences, 18th Edition, (1990), Mack Publishing Co., Easton, PA.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal  
10 administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

Also included are solid form preparations which are intended to be converted,  
15 shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type  
20 as are conventional in the art for this purpose.

Preferably the compound is administered orally.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve  
25 the desired purpose.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 1 mg to about 150 mg, preferably from about 1 mg to about 75 mg, more preferably from about 1 mg to about 50 mg, according to the particular application.

30 The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For

- 131 -

convenience, the total daily dosage may be divided and administered in portions during the day as required.

The amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration can range from about 1 mg/day to about 300 mg/day, preferably 1 mg/day to 75 mg/day, in two to four divided doses.


The methods of this invention described above using a compound of Formula I also include the use of one or more compounds of Formula I, and the methods of this invention described above using a compound of Formula I in combination with an H<sub>1</sub> receptor antagonist also include the use of one or more compounds of Formula I in combination with one or more H<sub>1</sub> receptor antagonists.

While the present has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

- 132 -

**WHAT IS CLAIMED IS:**

1. A compound of the formula:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

(1)  $R^1$  is selected from:

- (a) aryl;
- (b) heteroaryl;
- (c) heterocycloalkyl
- (d) alkyl;
- (e) cycloalkyl; or
- (f) alkylaryl;



wherein said  $R^1$  groups are optionally substituted with 1 to 4 substituents

independently selected from:

- (1) halogen;
- (2) hydroxyl;
- (3) lower alkoxy;
- (4)  $-CF_3$ ;
- (5)  $CF_3O-$ ;
- (6)  $-NR^4R^5$ ;
- (7) phenyl;
- (8)  $-NO_2$ ;
- (9)  $-CO_2R^4$ ;
- (10)  $-CON(R^4)_2$  wherein each  $R^4$  is the same or different;
- (11)  $-S(O)_mN(R^{20})_2$  wherein each  $R^{20}$  is the same or different H or alkyl group, preferably  $C_1$  to  $C_4$  alkyl, most preferably  $C_1$ - $C_2$  alkyl, and more preferably methyl;
- (12)  $-CN$ ; or

- 133 -

(13) alkyl; or

(2)  $R^1$  and X taken together form a group selected from:(3) X is selected from:  $=C(O)$ ,  $=C(NOR^3)$ ,  $=C(NNR^4R^5)$ ,(4)  $M^1$  is carbon;(5)  $M^2$  is selected from C or N;(6)  $M^3$  and  $M^4$  are independently selected from C or N;(7) Y is selected from: is  $-CH_2-$ ,  $=C(O)$ ,  $=C(NOR^{20})$  (wherein  $R^{20}$  is asdefined above), or  $=C(S)$ ;(8) Z is a  $C_1 - C_6$  alkyl group;


(9)  $R^2$  is a five or six-membered heteroaryl ring, said six-membered heteroaryl ring comprising 1 or 2 nitrogen atoms with the remaining ring atoms being carbon, and said five-membered heteroaryl ring containing 1 or 2 heteroatoms selected from: nitrogen, oxygen, or sulfur with the remaining ring atoms being carbon; said five or six membered heteroaryl rings being optionally substituted with 1 to 3 substituents independently selected from: halogen, hydroxyl, lower alkyl, lower alkoxy,  $-CF_3$ ,  $CF_3O-$ ,  $-NR^4R^5$ , phenyl,  $-NO_2$ ,  $-CO_2R^4$ ,  $-CON(R^4)_2$  wherein each  $R^4$  is the same or different,  $-CH_2NR^4R^5$ ,  $-(N)C(NR^4R^5)_2$ , or  $-CN$ ;

(10)  $R^3$  is selected from:

- (a) hydrogen;
- (b)  $C_1 - C_6$  alkyl;
- (c) aryl;
- (d) heteroaryl;
- (e) heterocycloalkyl;
- (f) arylalkyl;
- (g)  $-(CH_2)_e-C(O)N(R^4)_2$  wherein each  $R^4$  is the same or different,
- (h)  $-(CH_2)_e-C(O)OR^4$ ;

- 134 -

(i)  $-(CH_2)_e-C(O)R^{30}$  wherein  $R^{30}$  is a heterocycloalkyl group, such as, for example, morpholinyl, piperidinyl, piperazinyl or pyrrolidinyl, including



(j)  $-CF_3$ ; or

5 (k)  $-CH_2CF_3$ ;

wherein said aryl, heteroaryl, heterocycloalkyl, and the aryl portion of said arylalkyl are optionally substituted with 1 to 3 substituents selected from: halogen,  $-OH$ ,  $-OCF_3$ ,  $-CF_3$ ,  $-CN$ ,  $-N(R^{45})_2$ ,  $-CO_2R^{45}$ , or  $-C(O)N(R^{45})_2$ , wherein each  $R^{45}$  is independently selected from: H, alkyl, alkylaryl, or alkylaryl wherein said aryl moiety is substituted  
10 with 1 to 3 substituents independently selected from  $-CF_3$ ,  $-OH$ , halogen, alkyl,  $-NO_2$ , or  $-CN$ ;

(11)  $R^4$  is selected from: hydrogen,  $C_1 - C_6$  alkyl, aryl, alkylaryl, said aryl and alkylaryl groups being optionally substituted with 1 to 3 substituents selected from: halogen,  $-CF_3$ ,  $-OCF_3$ ,  $-OH$ ,  $-N(R^{45})_2$ ,  $-CO_2R^{45}$ ,  $-C(O)N(R^{45})_2$ , or  $-CN$ ; wherein  $R^{45}$  is  
15 as defined above;

(12)  $R^5$  is selected from: hydrogen,  $C_1 - C_6$  alkyl,  $-C(O)R^4$ ,  $-C(O)_2R^4$ , or  $-C(O)N(R^4)_2$  wherein each  $R^4$  is independently selected, and  $R^4$  is as defined above;

(13) or  $R^4$  and  $R^5$  taken together with the nitrogen atom to which they are bound forms a five or six membered heterocycloalkyl ring;

20 (14)  $R^6$  is selected from: alkyl, aryl, alkylaryl, halogen, hydroxyl, lower alkoxy,  $-CF_3$ ,  $CF_3O-$ ,  $-NR^4R^5$ , phenyl,  $-NO_2$ ,  $-CO_2R^4$ ,  $-CON(R^4)_2$  wherein each  $R^4$  is the same or different, or  $-CN$ ;

(15)  $R^{12}$  is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

(16)  $R^{13}$  is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

25 (17) a is 0 to 2;

(18) b is 0 to 2;

(19) c is 0 to 2;

(20) e is 0 to 5;

(21) m is 1 or 2;


30 (22) n is 1, 2 or 3; and

- 135 -

(23) p is 1, 2 or 3, with the proviso that when M<sup>3</sup> and M<sup>4</sup> are both nitrogen, then p is 2 or 3.


2. The compound of Claim 1 wherein R<sup>1</sup> is selected from:

- 5 (A) aryl;  
 (B) substituted aryl, wherein the substituents on said substituted aryl are selected from: (1) halo; or (2) alkyl; or (3) substituted alkyl;  
 (C) heteroaryl;  
 (D) substituted heteroaryl; or  
 10 (E) when R<sup>1</sup> is taken together with X, then the moiety is



3. The compound of Claim 2 wherein R<sup>1</sup> is selected from:

- 15 (A) phenyl;  
 (B) substituted phenyl wherein the substituents on said substituted phenyl are selected from: (1) halo; (2) alkyl; (3) alkyl substituted with halo;  
 (C) heteroaryl selected from: pyridyl, thienyl, pyrimidinyl, thiazolyl or pyridyl N-Oxide;  
 (D) alkyl substituted thiazolyl; or  
 20 (E) when R<sup>1</sup> is taken together with X, then the moiety is




wherein c is 0 or 1, and when c is 1 then R<sup>6</sup> is halo.

4. The compound of Claim 3 wherein R<sup>1</sup> is selected from:


- 25 (A) phenyl;  
 (B) substituted phenyl, wherein the substituents on said substituted phenyl are independently selected from: chloro, fluoro or trifluoromethyl;  
 (C) heteroaryl selected from:




- 136 -



(D) substituted heteroaryl of the formula:



(E) when  $R^1$  is taken together with X, then the moiety is



wherein c is 0 or 1, and when c is 1 then  $R^6$  is fluoro.


5. The compound of Claim 1 wherein  $R^1$  is selected from:

(A) phenyl;


(B) substituted phenyl, wherein the substituents on said substituted phenyl are independently selected from: chloro, fluoro or trifluoromethyl;

(C) pyridyl; or

(D) substituted heteroaryl of the formula:



(E) when  $R^1$  is taken together with X, then the moiety is




wherein c is 0 or 1, and when c is 1 then  $R^6$  is fluoro.

6. The compound of Claim 5 wherein  $R^1$  is pyridyl.

- 137 -

7. The compound of Claim 6 wherein  $R^1$  is



8. The compound of Claim 1 wherein X is  $=C(NOR^3)$ , and  $R^3$  is selected from H or alkyl.


9. The compound of Claim 8 wherein  $R^3$  is selected from H, methyl or ethyl.

10. The compound of Claim 9 wherein  $R^3$  is methyl.

11. The compound of claim 1 wherein: (1)  $M^2$  is nitrogen; and (2)  $M^3$  and  $M^4$  are selected such that: (a) one is carbon and the other is nitrogen, or (b) both are nitrogen.

12. The compound of Claim 11 wherein:  
 $M^3$  is carbon, and  $M^4$  is nitrogen;  
 $n$  is 2;  
 $a$  is 0 or 1;  
 $b$  is 0 or 1;  
 $c$  is 0 or 1, and when  $c$  is 1 then  $R^6$  is halo;  
 $e$  is 1 to 5; and  
 $p$  is 2.  
 $Y$  is  $=C(O)$ ; and  
 $Z$  is  $C_1$  to  $C_3$  alkyl.

13. The compound of Claim 1 wherein Z is



- 138 -


14. The compound of Claim 1 wherein  $R^2$  is a six membered heteroaryl ring.

15. The compound of Claim 14 wherein  $R^2$  is selected from pyridyl, pyridyl substituted with  $-NR^4R^5$ , pyrimidinyl, or pyrimidinyl substituted with  $-NR^4R^5$ .

5

16. The compound of Claim 15 wherein  $R^2$  is pyridyl substituted with  $-NH_2$ , or pyrimidinyl substituted with  $-NH_2$ .

17. The compound of Claim 16 wherein  $R^2$  is



10

18. The compound of Claim 1 wherein  $R^4$  is H or lower alkyl;  $R^5$  is H,  $C_1$  to  $C_6$ alkyl, or  $-C(O)R^4$ ;  $R^{12}$  is alkyl, hydroxy or fluoro; and  $R^{13}$  is alkyl, hydroxy or fluoro.

15

19. The compound of Claim 18 wherein  $R^4$  is H or methyl;  $R^5$  is H or methyl;  $R^{12}$  is hydroxy or fluoro; and  $R^{13}$  is hydroxy or fluoro.

20. The compound of Claim 1 wherein:

(1)  $R^1$  is selected from:

20

(A) aryl;

(B) substituted aryl, wherein the substituents on said substituted aryl


are selected from: (1) halo; or (2) alkyl; or (3) substituted alkyl;

(C) heteroaryl; or

(D) substituted heteroaryl; or

25

(E) when  $R^1$  is taken together with X, then the moiety is




(2) X is  $=C(NOR^3)$ ;

- 139 -

- (3)  $R^3$  is selected from H or alkyl;  
 (4)  $M^2$  is nitrogen;  
 (5) Y is  $=C(O)$ ;  
 (6)  $M^3$  and  $M^4$  are selected such that: (1) one is carbon and the other is  
 5 nitrogen, or (2) both are nitrogen;  
 (7) Z is  $C_1$  to  $C_3$  alkyl; and  
 (8)  $R^2$  is a six membered heteroaryl ring.

21. The compound of Claim 16 wherein:

- 10 (1)  $R^1$  is selected from:  
 (A) phenyl;  
 (B) substituted phenyl wherein the substituents on said substituted phenyl are selected from: (1) halo; (2) alkyl; (3) alkyl substituted with halo;  
 (C) heteroaryl selected from: pyridyl, thienyl, pyrimidinyl, thiazolyl or  
 15 pyridyl N-Oxide; or  
 (D) alkyl substituted thiazolyl; or  
 (E) when  $R^1$  is taken together with X, then the moiety is




wherein c is 0 or 1, and when c is 1 then  $R^6$  is halo;


- 20 (2)  $R^3$  is selected from H, methyl or ethyl;  
 (3) n is 2,  
 (4) a is 0 or 1,  
 (5) b is 0 or 1,  
 (6) c is 0 or 1 and when c is 1 then  $R^6$  is halo,  
 25 (7) e is 1 to 5,  
 (8) p is 2,  
 (9)  $R^4$  is H or lower alkyl,  
 (10)  $R^5$  is H,  $C_1$  to  $C_6$ alkyl, or  $-C(O)R^4$ ;  
 (11)  $R^{12}$  is alkyl, hydroxy or fluoro, and  
 30 (12)  $R^{13}$  is alkyl, hydroxy or fluoro.

- 140 -

22. The compound of Claim 21 wherein R<sup>2</sup> is



R<sup>1</sup> is




5

M<sup>2</sup> is nitrogen, M<sup>3</sup> is carbon, and M<sup>4</sup> is nitrogen.

23. The compound of Claim 1 selected from: Compound 23, 30, 31, 32, 33,  
 10 41, 44, 45, 49, 50, 52, 53, 54, 55, 56, 57A, 59, 65, 75, 76, 80, 82, 83, 88, 92, 99, 104,  
 105, 110, 111, 117, 121, 123, 127, 128, 200-241, 244-273, 275, 278-282, 287, 296,  
 301-439, or 446.


24. The compound of Claim 1 selected from: Compound 32, 54, 55, 253A,  
 15 287 or 320.

25. The compound of Claim 1 having the formula:




- 141 -

26. The compound of Claim 1 having the formula:




27. The compound of Claim 1 having the formula:




5

28. The compound of Claim 1 having the formula:




10

29. The compound of Claim 1 having the formula:




- 142 -

30. The compound of Claim 1 having the formula:




31. The compound of Claim 1 having the formula:



5

32. The compound of Claim 1 having the formula:



10

33. A pharmaceutical composition comprising: (a) an effective amount of a compound of any of Claims 1 to 32 and a pharmaceutically effective carrier; or (b) an effective amount of a compound of any of Claims 1 to 32, and an effective amount of H<sub>1</sub> receptor antagonist, and a pharmaceutically effective carrier.

15

34. A use of a compound of any of claims 1 to 32 for the manufacture of a medicament for treating: allergy, allergy-induced airway responses, congestion, cardiovascular disease, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the

- 143 -

central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine.

35. The use of Claim 34 wherein allergy-induced airway responses are  
5 treated.

36. The method of Claim 34 wherein allergy or nasal congestion is treated.

37. The use of a compound of any of Claims 1 to 32 and the use of an H<sub>1</sub>  
10 receptor antagonist for the manufacture of a medicament for treating: allergy, allergy-induced airway responses, and congestion.

38. The use of Claim 37 wherein said H<sub>1</sub> receptor antagonist is selected  
from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine,  
15 chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pynilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or  
20 triprolidine.

39. The use of Claim 38 wherein said H<sub>1</sub> receptor antagonist is selected  
from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

40. The use of Claim 39 wherein said H<sub>1</sub> receptor antagonist is selected  
25 from: loratadine or descarboethoxyloratadine.